

Maturation *in vitro* of immature human oocytes for clinical use

Kwang-Yul Cha and Ri-Cheng Chian¹

Infertility Medical Center, CHA General Hospital, College of Medicine, Pochon CHA University, Seoul, Korea

TABLE OF CONTENTS

Introduction	103
Regulation of human oocyte maturation	103
Utilization of immature human oocytes	108
Conclusions	115
Acknowledgements	115
References	115

Human oocyte maturation is considered as the reinitiation and completion of the first meiotic division from the germinal vesicle stage (prophase I) to metaphase II, and the accompanying cytoplasmic maturation for fertilization and early embryonic development. Immature human oocytes obtained from patients undergoing gynaecological surgery, or ovulation induction or having polycystic ovary syndrome (PCOS) can be matured and fertilized *in vitro*. To date, 80% of immature oocytes matured to metaphase II when cultured in maturation medium supplemented with gonadotrophins and 85% of matured oocytes fertilized and cleaved *in vitro*. Following transfer of these embryos, pregnancies and live births have been achieved. However, the capacity for oocyte maturation was different when the immature oocytes were retrieved from PCOS patients and when the oocytes were cryopreserved at germinal vesicle stage.

Key words: fertilization/germinal vesicle/human/maturation/oocyte

Introduction

Since the first successful human pregnancy from in-vitro fertilization (IVF) was achieved (Steptoe and Edwards, 1978), assisted reproductive technology has become the frontier of both infertility treatment and research. The core of an assisted reproduction programme is oocyte quality

because one of the major problems encountered by IVF clinics is to recognize the maturation state of oocytes obtained from stimulated or unstimulated follicles. The cytoplasm of the oocyte is of key interest in oocyte maturation. The best way to improve embryo quality is to improve oocyte quality. It is known that insufficient cytoplasmic maturation of the oocyte will fail to promote male pronuclear formation and will thus increase chromosomal abnormalities after fertilization (Thibault *et al.*, 1975). Successful fertilization, development and pregnancy have been achieved with immature human oocytes matured *in vitro* in some infertility centres (Veeck *et al.*, 1983; Prins *et al.*, 1987; Cha *et al.*, 1989, 1992, 1996; Paulson *et al.*, 1994; Trounson *et al.*, 1994; Barnes *et al.*, 1995, 1996; Nagy *et al.*, 1996; Edirisinghe *et al.*, 1997; Jaroudi *et al.*, 1997; Liu *et al.*, 1997; Russell *et al.*, 1997); however, little is known about what conditions are suitable for in-vitro maturation of human oocytes, enabling IVF and embryo development in culture. Although our understanding is currently incomplete, it is now clear that addition of gonadotrophins [follicle stimulating hormone (FSH), luteinizing hormone (LH)] to the culture medium is beneficial to cytoplasmic maturation *in vitro*. In this paper, we will review research and clinical applications utilizing immature human oocytes.

Regulation of human oocyte maturation

Human oocytes usually become arrested in prophase I of meiosis during fetal life. At birth, the oocytes remain in the dictyate phase and each ovary has ~500 000 healthy non-growing or primordial follicles (Baker, 1963). Throughout the reproductive life of the woman, cohorts of oocytes are removed from this non-growing pool and commence growth. Near the completion of growth, oocytes acquire the ability to reinitiate meiosis (Pincus and Enzmann, 1935). Following resumption of meiosis, the

¹To whom correspondence should be addressed at: McGill Reproductive Center, Department of Obstetrics and Gynecology, Royal Victoria Hospital, McGill University, Montreal, Quebec, Canada H3A 1A1. Fax: 514-843-1496. E-mail: rchian@rvhob2.lan.mcgill.ca

nuclear membrane dissolves (germinal vesicle breakdown; GVBD) and chromosomes progress from metaphase I to telophase I. The completion of the first meiotic division is characterized by the extrusion of the first polar body and formation of the secondary oocyte, both of which contain a haploid chromosome complement. The second meiotic division is initiated rapidly after completion of the first meiotic division and the oocytes reach metaphase II prior to ovulation. Oocyte maturation is defined as the reinitiation and completion of the first meiotic division from the germinal vesicle stage to metaphase II, with accompanying cytoplasmic maturation necessary for fertilization and early embryonic development. Most knowledge of the regulation of oocyte maturation comes from animal experiments. The reader should keep this in mind although similar concepts may be relevant to human oocytes.

Nuclear maturation

GVBD is initiated by either the pre-ovulatory surge of gonadotrophins (LH) or the atretic degeneration of the follicle *in vivo*. The competence to undergo GVBD was previously thought to be acquired during follicle growth. However, it has been reported that the acquisition of GVBD competence is not dependent on oocyte growth (Canipari *et al.*, 1984). Therefore, the acquisition of the competence to undergo GVBD could be a multi-step process. In humans, some germinal vesicle oocytes collected from both stimulated and unstimulated ovaries fail to undergo GVBD during culture *in vitro*. It seems that only the competence to undergo GVBD is only acquired during follicle growth, suggesting that an inherent capability for GVBD is still important.

Many potential factors mediate the cumulus cells' control of GVBD (Figure 1). High levels of cyclic adenosine monophosphate (cAMP) and purine hypoxanthine in the culture medium prevent oocyte GVBD (Heller *et al.*, 1981; Downs *et al.*, 1985; Eppig *et al.*, 1985; Hillensjö *et al.*, 1985; Törnell and Hillensjö, 1993). The oocyte and cumulus cells are coupled by gap junctions, suggesting that granulosa cells control GVBD via the cumulus cells. The gap junctions permit regulatory molecules, such as steroids, Ca^{2+} , inositol 1,4,5-trisphosphate (IP_3), cAMP and purines, to pass freely between the cytoplasm of the oocyte and cumulus cells (Dekel *et al.*, 1981). Addition of luteinizing hormone (LH) to the culture medium induces GVBD. LH probably induces GVBD by an indirect action mediated by cumulus cells because there are no LH receptors on oocytes (Dekel, 1988). The mechanism involves LH inducing the loss of communication between oocyte and cumulus cell complex,

thus terminating the flow of these substances into the oocyte (Berridge and Irvine, 1984; Larsen *et al.*, 1987; Downs *et al.*, 1988; Fagbohun and Downs, 1991). LH-induced GVBD may be also mediated by the $\text{IP}_3/\text{Ca}^{2+}$ pathway (for review see Eppig, 1993). It has been suggested that the hormone signalling mediate Ca^{2+} transients may be important at the time of GVBD (for review see Whitaker, 1996).

Results from animal experiments indicate that protein synthesis is required for GVBD (Stern *et al.*, 1972; Schultz and Wassarman, 1977; Sirard and First, 1988; Sirard *et al.*, 1989). It seems that protein synthesis is also needed for GVBD of human oocytes because protein synthetic patterns differ before and after GVBD (Gifford *et al.*, 1987; Schultz *et al.*, 1988). After GVBD, not all oocytes can mature and extrude the first polar body. The competence to undergo GVBD and to progress from metaphase I to metaphase II is acquired separately during culture *in vitro* (Szybek, 1972; Iwamatsu and Yanagimachi, 1975; Sorensen and Wassarman, 1976). Some human oocytes undergo GVBD when cultured *in vitro*, but become arrested at metaphase I. Thus, the competence to undergo GVBD and competence to progress through metaphase I to metaphase II are separately and sequentially acquired. It is not clear, however, what causes oocytes to complete maturation from metaphase I to metaphase II. It is well known that the cytoplasmic maturation during this period is very important in order to support early development. Protein synthesis is needed for the progression of oocytes from the germinal vesicle stage to metaphase II, as well as for the maintenance of the metaphase II arrest (Clarke and Masui, 1983; Gerhart *et al.*, 1984).

Inhibition of protein synthesis in oocytes results in failure to activate maturation promoting factor (MPF) activity (Masui and Clarke, 1979; Maller, 1985; Hashimoto and Kishimoto, 1988). Cytoplasmic proteins, MPF and cytosolic factor (CSF; Shibuya and Masui, 1989) regulate oocyte nuclear maturation. Molecular characterization of MPF has shown that active MPF is a protein dimer composed of catalytic p34^{cdc2} serine/threonine kinase, and regulatory cyclin B subunits (Dunphy *et al.*, 1988; Gautier *et al.*, 1988; 1990; Pines and Hunter, 1989). The p34^{cdc2} serine/threonine kinase is the product of the *cdc2* gene, first identified in fission yeast. The p34^{cdc2} -cyclin heterodimer, a protein kinase, has four phosphorylation sites that are regulated by kinase and phosphatase activities. Activation of MPF requires dephosphorylation at Thr 14 and Tyr 15 and phosphorylation of Thr 161 (for review see Piwnicka-Worms *et al.*, 1993). Kinases and phosphatases capable of modifying p34^{cdc2} have been identified by genetic analysis in yeast




Figure 1. Hypothetical model for the possible participation of factors in germinal vesicle breakdown (GVBD) of human oocytes. LH = luteinizing hormone; GV = germinal vesicle; MPF = maturation promoting factor; CSF = cytotstatic factor; IP₃ = inositol 1,4,5-trisphosphate.

(wee1, cdc25) and in frog oocytes (cak: cdc activating kinase) (for review see Whitaker, 1996).

G2/M arrest always finds the oocyte with a high content of inactive p34^{cdc2} complex in its tyrosine-phosphorylated form. It induces GVBD when p34^{cdc2}-cyclin is activated (Ookata *et al.*, 1992). Before fertilization, human oocytes arrest firmly at metaphase II. At this point, MPF kinase activity is high and cyclin destruction has been inhibited. This is due to CSF. It is known that mos protein, the product of the *c-mos* proto-oncogene, has the same effect as CSF in arresting mitosis at metaphase with high p34^{cdc2} activity. Clearly, meiotic arrest is usually explained in terms of activation and inactivation of cell cycle control proteins, particularly p34^{cdc2}. Oocytes need p34^{cdc2} activation to enter meiotic metaphase and p34^{cdc2} inactivation to proceed through anaphase into the next cell cycle. Therefore, the metaphase II arrest is due to the transcription of *c-mos* as the oocyte matures.

The *c-mos* gene plays a critical role in regulation of MPF activity during meiosis. The product of *c-mos* is expressed early in oocyte maturation and disappears immediately after fertilization (Sagata *et al.*, 1989; Watanabe *et al.*, 1989). *c-mos* encodes a protein-serine/threonine kinase known as Mos. Mos has been proposed to enhance MPF activity via several mechanisms (Figure 2). Direct activation of MPF occurs through phosphorylation of the cyclin-B subunit of MPF (Roy *et al.*, 1990), or indirectly via mitogen-activated protein kinase (MAP kinase) (Nebreda and Hunter, 1993). In addition, Mos may inhibit proteolytic degradation of cyclin-B between meiosis I and

meiosis II (O'Keefe *et al.*, 1991), and thus maintain high MPF activity (Kubiak *et al.*, 1993) because Mos is required for cyclin B accumulation and this newly synthesized protein is required to maintain MPF activity during meiosis of oocytes (O'Keefe *et al.*, 1989; Minshull *et al.*, 1991). Thus, inhibition of protein synthesis will block the effects of Mos activity. The *c-mos* gene is an abundant maternal message in the oocytes. Expression of *c-mos* has recently been reported in human oocytes (Goldman *et al.*, 1987; Mutter and Wolgemuth, 1987; Pal *et al.*, 1994), suggesting that Mos may be a critical regulator of human oocyte meiosis. Surprisingly, using reverse transcriptase-polymerase chain reaction (RT-PCR), Li *et al.* (1993) detected *c-mos* mRNA in various human cell lines, suggesting a wider role for *c-mos* kinase.

Like Mos, MAP kinase was first identified in somatic cells and has now been revealed as central to the regulation of meiotic arrest in oocytes. MAP kinase is also a serine/threonine kinase but is activated, not inhibited, by tyrosine phosphorylation. It is known that MAP kinase is activated during oocyte maturation, that it regulates meiosis and has CSF activity. Activation of MAP kinase precedes activation of p34^{cdc2}. Blocking MAP kinase activity prevents GVBD. Mos protein stimulates MAP kinase activity, but does not activate p34^{cdc2} (Nebreda and Hunter, 1993). The time course of MAP kinase activation in oocytes parallels that of *c-mos* transcription (Sagata *et al.*, 1988; Ferrell *et al.*, 1991); therefore, it seems that the MAP kinase activity in oocytes is due to Mos. The phosphorylation cascade of Mos and MAP kinase may play an important role




Figure 2. The regulatory pathways surrounding maturation promoting factor (MPF). p34^{cdc2} combines with its activating subunit, cyclin B. The p34^{cdc2}-cyclin B heterodimer, a protein kinase, has four phosphorylation sites that are regulated by kinase and phosphatase activities. The phosphorylation cascade of Mos and MAP kinase plays an important role in MPF activity for germinal vesicle breakdown (GVBD).

in meiotic and mitotic cell cycles. Mos activates MAP kinase to arrest oocytes at metaphase II; however, the connection between Mos/MAP kinase and p34^{cdc2} activation at a molecular level is not clear. After fertilization, Mos activity is lost (Lorca *et al.*, 1993). Cyclin disappears before Mos (Kubiak *et al.*, 1993). The evidence indicates that calcium is the signal to break the metaphase II arrest of oocytes (for review see Swann, 1996).

Cyclin was originally identified in sea urchin oocytes as a protein whose level greatly increased upon fertilization and subsequently oscillated during the early cell divisions of the embryo (Evans *et al.*, 1983). The level of cyclin increases steadily through interphase, peaks at the G2/M phase transition, and falls precipitously at each mitosis. Cyclins have been divided into three classes, G1, A and B, based on their amino acid similarity and the timing of their appearance during the cell cycle (Minshull *et al.*, 1989; Hunter, 1991; Pines and Hunter, 1991). Two isoforms of cyclin-B have been described in the mouse (Chapman and Wolgemuth, 1992, 1993). The expression patterns of cyclin-B1 and -B2 differ, with the cyclin-B1 isoform predominantly expressed in the oocytes. Cyclin B is also phosphorylated and dephosphorylated during oocyte maturation (Whitaker and Patel, 1990). Cyclin-B1 was expressed in oocytes, embryos and granulosa cells from both the human and monkey (Heikinheimo *et al.*, 1995, 1996), indicating that the expression patterns of *c-mos*, cyclin-B1 and β -actin are important for proper oocyte development in human oocytes and preimplantation embryos.

Cytoplasmic maturation

During oocyte growth and maturation, products are synthesized and stored. These products support development after fertilization until the embryonic genome becomes transcriptionally active and embryo-derived messages begin to regulate embryogenesis. Most of the RNA present in mammalian oocytes is synthesized and accumulated during the period of oocyte growth (Bachvarova, 1974; Jahn *et al.*, 1976; Bachvarova and DeLeon, 1980; Sternlicht and Schultz, 1981; Piko and Clegg, 1982). RNA synthesis continues at a low level to within 1 h of GVBD and some of the newly synthesized RNA is released into the cytoplasm before GVBD (Bloom and Mukherjee, 1972; Rodman and Bachvarova, 1976; Wassarman and Letourneau, 1976). It is known that the metabolism of the oocyte is characterized by active transcription and translation during the pre-ovulatory period (Telford *et al.*, 1990; Wassarman and Kinloch, 1992); however, transcription ceases at the time of ovulation, so the oocyte and early pre-embryo are dependent upon the pool of mRNA and protein accumulated during the pre-ovulatory period (Telford *et al.*, 1990).

The quality of the oocyte predominantly depends upon the maturity of the ooplasm. The cytoplasm prepares for fertilization and early embryonic development in parallel with nuclear maturation. If cytoplasmic maturation does not occur properly, the oocyte will fail to fertilize and develop successfully (Thibault *et al.*, 1975; Thibault, 1977). Mammalian oocytes are surrounded by a compact cumulus

cell layer during folliculogenesis. Cumulus cells respond to gonadotrophins and are known to secrete various substances. These substances not only control nuclear maturation but also perform an important role in cytoplasmic maturation. Beneficial effects of cumulus cells on early development have been reported for rabbit (Robertson and Baker, 1969), mouse (Cross and Brinster, 1970), rat (Vanderhyden and Armstrong, 1989), human (Kennedy and Donahue, 1969; Gregory and Leese, 1996), ovine (Staigmiller and Moor, 1984), and bovine (Sirard *et al.*, 1988; Fukui, 1990; Chian *et al.*, 1994) oocytes. Besides cytoplasmic changes involved in the control and direction of meiotic progression, other important cytoplasmic changes occur during oocyte maturation (Mattioli 1992). In pigs, the formation of a male pronucleus within a fertilized oocyte depends on the presence of cumulus cells during oocyte maturation (Mattioli *et al.*, 1988). The events of male pronucleus formation after sperm penetration are also affected by the presence or absence of cumulus cells during bovine oocyte maturation *in vitro* (Chian *et al.*, 1994). These results suggest that the ability for male pronuclear formation is acquired during late folliculogenesis. The protein synthesis pattern is different between oocytes with and without cumulus cells and FSH modulates the protein synthesis pattern of cumulus cell-intact oocytes (Chian and Sirard, 1995).

The cumulus cells are distinct from the mural granulosa cells. The cumulus cells have different levels of progesterone secretion and different responses to gonadotrophins (Dirnfeld *et al.*, 1993; Goldman *et al.*, 1993). The cumulus cells also differ in the expression of specific mRNA in comparison with the mural granulosa cells (Braw-Tal, 1994). Several regulatory factor transcripts have been demonstrated in human oocytes by RT-PCR, including oestrogen receptor, *c-mos* and *c-raf* mRNA (Wu *et al.*, 1993; Wu and Wolgemuth, 1995). Therefore, these regulatory factors may be involved in the events of cytoplasmic maturation of the oocyte. Thyroid hormone, tri-iodothyronine (T_3), synergizes with FSH to increase LH receptors and to increase progesterone secretion by granulosa cells from porcine follicles (Maruo *et al.*, 1987, 1992). A recent report indicated that T_3 may also be important for oocyte maturation since four isoforms of thyroid hormone receptor (TR) mRNA (TR- α -1, TR β -1, TR β -2 and c-erbA α -2) are expressed in the human oocyte, and the granulosa cells as well as on the cumulus cells (Zhang *et al.*, 1997). These results suggest that thyroid hormone may both have direct effects on the oocytes, and indirect effects via the cumulus cells. The expression of TR mRNAs in the oocyte, cumulus cells and granulosa cells raises the question of whether addition of T_3 to the maturation medium would improve maturation *in vitro* of

immature human oocytes in a more rapid or normal manner (Zhang *et al.*, 1997).

It is also known that different culture conditions for oocyte maturation *in vitro* affect fertilization and embryonic development. Normally, FSH is added to the culture medium for oocyte maturation *in vitro*, but addition of FSH does not increase the nuclear maturation of rat (Vanderhyden and Armstrong, 1989), monkey (Morgan *et al.*, 1991) and human (Durinzi *et al.*, 1997) oocytes and FSH initially has an inhibitory action on mouse oocyte maturation (Downs *et al.*, 1988). It has been reported that mouse oocyte GVBD is inhibited by FSH due to the rise in the levels of cAMP in cumulus cells (Eppig *et al.*, 1983; Schultz *et al.*, 1983).

FSH is important in the development of pre-ovulatory follicles. The enhanced FSH responsiveness of pre-ovulatory follicles appears to result from an increase in the content of the stimulatory G-protein of the adenyl cyclase system (for review see Richards and Hedin, 1988). FSH is necessary for the induction of LH receptors in pre-ovulatory follicles; however, the mechanism of FSH on nuclear and cytoplasmic maturation of oocytes is not clear. It was found that FSH levels in the follicular fluid of metaphase I and metaphase II oocytes were significantly higher than in follicles containing germinal vesicle oocytes in women undergoing oocyte retrieval for stimulated IVF cycles (Laufer *et al.*, 1984). The addition of FSH to oocyte culture medium does not significantly increase the ability of the oocyte to reach metaphase II (Durinzi *et al.*, 1997); however, maturation medium with FSH significantly increases fertilization (Schroeder *et al.*, 1988) and early embryo development (Jinno *et al.*, 1989; Morgan *et al.*, 1991). The ability of FSH to increase the developmental capacity of mouse oocytes maturing *in vitro* varies depending on the age and prior gonadotrophin priming *in vivo* (Eppig *et al.*, 1992).

Sperm penetration triggers the events that signal the oocyte has been activated and has entered the programme of embryonic development. The activated oocyte causes the exocytosis of the cortical granules, which prevents polyspermy and the completion of meiosis to start the first embryonic mitosis. These events of oocyte activation at fertilization are mediated by a sperm-induced increase in the concentration of intracellular free Ca^{2+} (Kline and Kline, 1992; Whitaker and Swann, 1993). The development of the ability of the oocyte to release Ca^{2+} in response to the fertilizing spermatozoa is an essential step during oocyte maturation (for review see Carroll *et al.*, 1996). Recent evidence has revealed that Ca^{2+} release mechanisms are modified during oocyte maturation (Chiba *et al.*, 1990; Fujiwara *et al.*, 1993; Mehlmann and Kline, 1994). When immature human oocytes were cultured *in vitro*, they acquired

the capacity to undergo a single large oscillation of intracellular Ca^{2+} ; however, subsequent sustained oscillations were not observed in some immature oocytes, indicating that these oocytes failed to develop fully competent Ca^{2+} signalling mechanisms during maturation *in vitro* (Herbert *et al.*, 1997). It seems that the maximal sensitivity of Ca^{2+} release is reached in the final stages of oocyte maturation, just before the optimal time for fertilization.

In oocytes, cytosolic free Ca^{2+} is increased by the release of Ca^{2+} from intracellular Ca^{2+} stores in the endoplasmic reticulum. The release of Ca^{2+} from the stores is controlled by two types of receptors/ Ca^{2+} release channels, one sensitive to IP_3 and the other to ryanodine (Lai *et al.*, 1988; Furuichi *et al.*, 1989). These two Ca^{2+} release channels are sensitive to cytosolic Ca^{2+} such that low concentrations stimulate opening while high concentrations inhibit the receptors (Bezprozvanny *et al.*, 1991; Iino and Endo, 1992). The mechanism of the Ca^{2+} oscillations that occur during fertilization of mammalian oocytes is still unclear. There is considerable evidence to indicate that IP_3 plays a primary role in generating Ca^{2+} signals in mammalian oocytes (Miyazaki *et al.*, 1992; Tesarik and Sousa, 1996; Berridge, 1996; Swann and Lawrence, 1996). It has been suggested that human oocyte activation at fertilization is mediated by the IP_3 Ca^{2+} release system because the Ca^{2+} oscillations were inhibited by caffeine (Herbert *et al.*, 1997). The development of Ca^{2+} release mechanism during oocyte maturation so that the fertilizing sperm initiates a Ca^{2+} signal sufficient to stimulate oocyte activation is still a mystery. Changes may be brought about by modifications in the sensitivity of Ca^{2+} release through the IP_3 receptors. Ca^{2+} release channels are also regulated by phosphorylation (Nakade *et al.*, 1994) and changes in redox state (Salama *et al.*, 1992) as well as by interactions with cytoplasmic proteins such as ankyrin (Bourguignon *et al.*, 1993) and FK506-binding protein (Brillantes *et al.*, 1994; Cameron *et al.*, 1995). Therefore, it is an important that the oocyte develops the ability to respond to spermatozoa with Ca^{2+} transients during cytoplasmic maturation.

Oocyte polarity and spatial patterning during maturation should also receive attention (for review see Gardner, 1996; Edwards and Beard, 1997). Microvilli are distributed evenly during human oocyte maturation (Pickering *et al.*, 1988) and a microvillus-free region was observed as a consequence of ageing in culture in human oocytes (Santello *et al.*, 1992). Several proteins display a polarized cortical distribution at different maturation stages of human oocytes (Ji *et al.*, 1996). The maternally encoded mRNA to establish a critical polarity and spatiality seems to be involved in oocyte growth and maturation. However, very little is known about the polarity and the spatiality of maternal mRNA in mammalian

oocytes, especially during oocyte maturation. Moreover, leptin, a 16 kDa protein product of the obese gene, and its receptor isoforms are in human pre-ovulatory follicles and in mature oocytes, suggesting that leptin may play some role in transcriptional regulation of the oocyte and early embryo development (Cioffi *et al.*, 1997).

It is only after activation of the embryonic genome that the embryo begins to be controlled by its own unique genome. The information required for the embryonic cells to reorganize their surface according to their special orientation within the embryo is probably part of the maternal genetic message inherited from the oocyte in the form of pre-synthesized RNA or proteins and utilized to guide developmental processes in the early embryo. In humans, this is thought to occur between the 4- and 8-cell stages (Tesarik *et al.*, 1986; Braude *et al.*, 1988). Early cleavage arrest has been demonstrated in most animals (Flach *et al.*, 1982; Davis, 1985; Crosby *et al.*, 1988; Frei *et al.*, 1989; Sakkas *et al.*, 1989). In human pre-embryos, the *c-mos* transcripts disappear rapidly and are not detectable beyond the 5-cell stage and expression of cyclin-B1 is somewhat paralleled and increased after the 6-cell stage (Heikinheimo *et al.*, 1995). This suggests that active transcription of these genes in human pre-embryos is from the 4-cell stage onwards. However, Artley *et al.* (1992) have examined the relationship between gene expression and cleavage arrest by investigating the patterns of protein synthesis in cleavage-arrested human pre-embryos cultured *in vitro*, indicating that cleavage arrest is not always accompanied by failure of activation of the genome. The underlying mechanisms responsible for cleavage arrest are unclear. Possible causes include inadequate culture conditions (Gandolfi and Moor, 1987; Menezo *et al.*, 1990), inherent or induced abnormality (Macas *et al.*, 1990; Pickering *et al.*, 1990) and failure of embryonic gene expression (Braude *et al.*, 1990). The optimal culture condition *in vitro* seems to promote the expression of the embryonic genome. It has been reported that transcription activity of mouse blastocysts can be enhanced by alterations in culture conditions (Ho *et al.*, 1994). Thus, it is important for effective gene transcription of the human pre-embryo that the culture conditions of assisted reproductive techniques are correct.

Utilization of immature human oocytes

Attempts to study human fertilization in culture were first reported by Rock and Menkin (1944). It was confirmed by Edwards (1965a,b) that human follicular oocytes are able to mature *in vitro* when they are isolated from follicles and placed in appropriate culture medium. Human oocytes surrounded by cumulus cells have a higher maturation rate

than those without these cells (Kennedy and Donahue, 1969). The first successful fertilization *in vitro* of human oocytes matured *in vitro* was reported by Edwards *et al.* (1969). Subsequently, the first successful human pregnancy from IVF of the oocytes retrieved from natural ovulatory cycles was achieved by Steptoe and Edwards (1978). Thereafter, the successful fertilization, development and pregnancy with immature human oocytes matured *in vitro* has been reported continuously (Veeck *et al.*, 1983; Prins *et al.*, 1987; Cha *et al.*, 1991, 1992, 1996; Paulson *et al.*, 1994; Trounson *et al.*, 1994; Barnes *et al.*, 1995, 1996; Nagy *et al.*, 1996; Edirisinghe *et al.*, 1997; Jaroudi *et al.*, 1997; Liu *et al.*, 1997; Russell *et al.*, 1997). However, very little is known about the maturation and developmental capacity of immature human oocytes for clinical application.

Oocytes from stimulated ovaries

For clinical application, most oocytes are obtained from ovarian follicles of women who have been treated with hormones to induce multiple follicular growth. The use of superovulation hormones has resulted in follicular asynchrony (Laufer *et al.*, 1984; Bomsel-Helmreich *et al.*, 1987). When oocytes were retrieved from stimulated ovaries before the LH surge, most of the oocytes were at the germinal vesicle stage (Bomsel-Helmreich *et al.*, 1987); however, oocytes retrieved after human chorionic gonadotrophin (HCG) administration are frequently maturing (metaphase I) and pre-ovulatory (metaphase II) oocytes from the cohort of developing follicles. At the same time, a few germinal vesicle oocytes may be retrieved, despite having been exposed to an ovulatory dose of HCG 36 h before aspiration. These oocytes are capable of undergoing spontaneous nuclear maturation *in vitro*, and then normal fertilization and development. Studies on oocytes from animals have shown that the endocrine environment plays an important role in ensuring normal cytoplasmic maturation with respect to subsequent fertilization (Moor and Trounson, 1977). The inclusion of human menopausal gonadotrophin (HMG) in the culture medium for immature human oocytes results in improved maturation and fertilization rates (Prins *et al.*, 1987; Table I). It has been reported that pregnancies have been obtained following the transfer of immature oocytes from stimulated cycles (Veeck *et al.*, 1983; Prins *et al.*, 1987; Nagy *et al.*, 1996; Liu *et al.*, 1997; Jaroudi *et al.*, 1997; Edirisinghe *et al.*, 1997).

In addition to the nuclear events, oocyte cytoplasmic maturation is necessary for normal fertilization and subsequent embryonic development. Human IVF programmes have indicated that delaying insemination improves fertilization rates, presumably by allowing the completion of cytoplasmic maturation for those oocytes

that have not completely matured at the time of retrieval (Trounson *et al.*, 1982). Increasing FSH levels within the follicle is coincident with the generation of a positive signal necessary to complete oocyte maturation in humans *in vivo*. This suggests that this signal may be linked to the dynamics of growth factors within the follicle itself (Gómez *et al.*, 1993a,b). Recently, it has been reported that LH is required during the peri-ovulatory period for normal embryo development *in vivo* (Weston *et al.*, 1996). Fertilization and cleavage rates were improved by the addition of 150 mIU/ml HMG into the medium if oocytes derived from different size of follicles were cultured *in vitro* shortly before insemination (Zhang *et al.*, 1993; Table II). This indicates that the presence of gonadotrophin during the short period of culture before insemination may also have provided the oocytes with a proper hormonal environment for both nuclear and cytoplasmic maturation.

Table I. Meiotic maturation and fertilization rates of immature human oocytes (germinal vesicle) derived from stimulated cycles^a

	Group I (no HMG)	Group II (+ HMG)
Total no. of oocytes	31	34
Oocytes matured (1st polar body) (%)	11 (35.6)	25 (73.5) ^b
Oocytes fertilized/inseminated oocytes(%)	4/11 (36.4)	16/25 (64.0) ^b
Oocytes fertilized/total oocytes (%)	4/31 (12.9)	16/34 (47.1) ^b

^aOocytes were cultured with Ham's F-10 medium supplemented with 7.5% maternal serum in the absence (group I) and presence (group II) of 75 mIU human menopausal gonadotrophin (HMG).

Data from Prins *et al.* (1987).

^bSignificantly different from group I.

Different ovulation induction protocols may result in variations in oocyte quality, fertilization and cleavage rates (Boué and Boué, 1973; De Sutter *et al.*, 1991a,b). Fertilization rates are different when using clomiphene citrate (CC)/HCG and gonadotrophin analogue (GnRHa) protocols (De Sutter *et al.*, 1992). GnRHa is now used routinely to prevent spontaneous gonadotrophin surges during ovarian stimulation for IVF programmes. The short and long GnRHa protocols are usually used and both regimens prevent premature LH surges. Although there is no significant difference with respect to clinical pregnancy rates per cycle between the long and short protocols (Frydman *et al.*, 1988; Garcia *et al.*, 1990; Maroulis *et al.*, 1991; Acharya *et al.*, 1992; Hughes *et al.*, 1992), the oocyte maturity, fertilization and cleavage rates are different (Greenblatt *et al.*, 1995). Furthermore, several studies have suggested that high concentrations of LH during the follicular phase of stimulation can have a negative effect on oocyte quality, pregnancy rate and the incidence of miscarriage (Stanger and

Yovich, 1985; Howles *et al.*, 1986; Homburg *et al.*, 1988; Regan *et al.*, 1990). HMG, which is a 1:1 mixture of FSH and LH, and some centres have substituted HMG partially or completely by pure FSH (Venturoli *et al.*, 1986). The highly purified FSH short-term stimulation protocol synchronizes oocyte maturation better than a comparable stimulation with HMG (Imthurn *et al.*, 1996). Moreover, increased fertilization rates have been reported with the addition of granulosa cells to culture medium used to mature oocytes with compact cumulus cells for 5–8 h after recovery in CC/HCG-stimulated protocols (Dandekar *et al.*, 1991). In addition, it must be noted that 18% of in-vitro maturing oocytes had gross meiotic aberrations (Racowsky and Kaufman, 1992). It is possible that these abnormalities also occur during maturation *in vivo* and after gonadotrophin stimulation of follicular growth for mature oocyte recovery. It is reported that aneuploidy occurs at high rates in mature oocytes (20–31%) recovered in stimulated cycles (Wramsby and Liedholm, 1984; Martin *et al.*, 1986; Bongso *et al.*, 1988; Djalali *et al.*, 1988; Papadopoulos *et al.*, 1988; Plachot *et al.*, 1988; Ma *et al.*, 1989; Pellestor, 1991; Gras *et al.*, 1992; Racowsky *et al.*, 1997).

Cellular, metabolic and cytoplasmic studies of mature human oocytes obtained from stimulated cycles provide explanations for some types of early developmental failures. Developmentally lethal chromosomal abnormalities are a common defect in mature human oocytes that may affect $\geq 25\%$ of normal-looking mature human oocytes retrieved from stimulated cycles (Plachot *et al.*, 1988; Tarin *et al.*, 1991; Delhanty and Handyside, 1995; Wall *et al.*, 1996). Specific cytoplasmic pathologies present in the oocyte at the time of insemination correlate with both the ability to fertilize and the ability of the fertilized oocyte to develop progressively (Van Blerkom, 1994). Metabolic differences between morphologically equivalent metaphase II oocytes retrieved from the same ovary have been observed and related to the developmental ability of the resulting embryo (Van Blerkom *et al.*, 1995; Barnett and Bavister, 1996; Van Blerkom, 1996). Very high frequencies of chromosomal aneuploidy have been observed in oocytes with cytoplasmic

defects, suggesting that these genetic abnormalities may develop during the pre-ovulatory stages in association with or as a result of degenerative cytoplasmic alterations (Van Blerkom and Henry, 1992). Some chromosomal anomalies are first detectable during early meiotic metaphase, therefore it is also possible that the damage of DNA may occur in earlier oogenesis (for review see Ashwood-Smith and Edwards, 1996). Increasing maternal age in naturally cycling women (Battaglia *et al.*, 1996a) and the unsuitable micro-environment for cytoplasmic maturation *in vivo* (Delhanty and Handyside, 1995), such as a lowering of intracellular pH (Van Blerkom, 1996) may be associated with high frequencies of spindle defects and chromosomal segregation disorders. Cytoplasmic maturation is also important to the development of microtubule organizing centres (MTOC) in the oocyte (Van Blerkom, 1991; Wickramasinghe *et al.*, 1991). A recent report indicated that the human oocyte develops MTOC as meiotic maturation proceeds beyond the prophase I arrest (Battaglia *et al.*, 1996b).

Anaesthesia might also affect oocyte maturation, fertilization and early cleavage *in vitro*. Normally, the procedure of ultrasound-guided transvaginal oocyte retrieval is carried out under either general, epidural or local anaesthesia. Propofol (diprivan) is one of the commonest medications used for anaesthesia in this situation. Propofol has been reported to accumulate in the follicular fluid (Palot *et al.*, 1988). Its concentrations in follicular fluid seem to increase steadily throughout the oocyte retrieval procedure, suggesting that Propofol may have deleterious effects on reproductive outcome following IVF (Coetsier *et al.*, 1992). Using a mouse IVF model, it has been suggested that Propofol is detrimental to oocyte maturation, fertilization and subsequent cleavage *in vitro* (Depypere *et al.*, 1991; Janssenswillen *et al.*, 1996; Alsalili *et al.*, 1997). However, it is unclear how Propofol affects oocyte maturation during culture *in vitro*. It is possible that the oocyte chromatin may be damaged by such anaesthetic agents since it is known that mammalian oocytes are unable to recognize damaged chromatin but will continue meiosis (Chian *et al.*, 1992; Pyrzynska *et al.*, 1996; Fulka, *et al.*, 1997).

Table II. Effect of human menopausal gonadotrophin (HMG) in maturation medium (MT6 medium +10% maternal serum) on in-vitro fertilization and cleavage of human oocytes derived from stimulated cycles^a

Follicle size (ml) ^b	No. of oocytes	Cumulus expanded	HMG (mIU)	Fertilization (%) ^c	Cleavage (%) ^c
32	44	3	0	10 (23)	4 (40)
32	51	5	150	25 (49)	19 (76)
>2	178	178	0	116 (65)	113 (97)

^aData from Zhang *et al.* (1993).

^bVolume of follicle fluid aspirated.

^cSignificantly different from each other ($P < 0.05$).




Figure 3. Human germinal vesicle oocytes retrieved from stimulated (a) and unstimulated ovaries (b). Both oocytes display compact cumulus cell layers. The intact germinal vesicle of the oocytes could be observed clearly under a microscope (original magnification $\times 400$).

Normally, 85–90% of the oocytes retrieved from stimulated cycles in IVF programmes are in metaphase II; however, 10–15% of oocytes derived from stimulated cycles are still immature (K.-Y. Cha *et al.*, unpublished data). Therefore, it is also important to improve the quality of these oocytes for clinical purposes. It has been reported that cumulus-free germinal vesicle oocytes retrieved from stimulated cycles can be matured *in vitro* by co-culturing with Vero cell monolayers (Janssenswillen *et al.*, 1995). The time course of oocyte GVBD and maturation were different between the oocytes retrieved from stimulated and unstimulated ovaries (Figure 3) and that the ability of the oocyte fertilization was also different between these two groups (R.-C. Chian *et al.*, unpublished data). The first GVBD was observed 6 h after culture *in vitro* in the oocytes retrieved from stimulated ovaries, but GVBD occurred 12–15 h after culture *in vitro* in the oocytes retrieved from unstimulated ovaries (Figure 4). However, the final percentages of GVBD were not different between the oocytes derived from stimulated (83.3%) and unstimulated (87.5%) cycles. GVBD occurred significantly earlier in the oocytes retrieved from stimulated ovaries than unstimulated ovaries. As discussed above, LH probably induces GVBD by an indirect action mediated by cumulus cells because there are no LH receptors on the oocytes (Dekel, 1988). Many factors may mediate the cumulus cells' control of GVBD; the mechanism involves LH inducing the loss of communication between oocyte and the cumulus cell complex, thus terminating the flow of some substances into the oocytes (Berridge and Irvine, 1984; Larsen *et al.*, 1987; Downs *et al.*, 1988; Fagbohun and Downs, 1991; Downs, 1995). From this




Figure 4. The time course of human oocyte germinal vesicle breakdown (GVBD) during culture *in vitro*. The final percentages of GVBD were not different between the oocytes retrieved from stimulated (—●—) and unstimulated (—○—) ovaries.

we may explain why GVBD occurs earlier in the oocytes derived from stimulated ovaries than from unstimulated ovaries, because the oocytes retrieved from stimulated ovaries have already been exposed to the action of LH/HCG stimulation in the follicles.

Extremely low fertilization rates are usually obtained after standard insemination of matured germinal vesicle oocytes from ovulation induction, suggesting that intracytoplasmic sperm injection (ICSI) is the best option, even where the sperm parameters are not impaired (Nagy *et al.*, 1996). It is not clear at the present time why the fertilization rates are so

low. It may be due to alterations of the zona pellucida (zona hardening) as a result of the oocyte maturation *in vitro* in these oocytes.

Oocytes from unstimulated ovaries

There are not many studies on immature human oocytes matured *in vitro* because of the limitation of materials. For clinical application and research, immature follicular oocytes can be obtained from consenting patients undergoing tuboplasty, Caesarean section, and oophorectomy. The method of collection of follicular immature oocytes from surgically removed ovaries were described previously by Cha *et al.* (1991). In the case of donors who underwent tuboplasty or Caesarean section, immature follicular oocytes were aspirated by puncturing visible follicles with a 21-gauge needle attached to a syringe filled with culture medium. The results indicate that the age, pathology, day of the menstrual cycle, and cyclic versus non-cyclic ovaries of donors, are factors influencing the number of immature oocytes (Cha, 1992). Immature oocytes from the luteal phase had a significantly higher maturation rate than those obtained during the follicular phase. The number of immature human oocytes collected from an ovary decreased as the patient's age increased; however, there was no statistical difference in the rate of oocyte maturation and cleavage among donors in the different age groups (Cha *et al.*, 1992).

Table III. Details of seven pregnancies using immature follicular oocytes (Cha, 1995)

Number	Maturation media ^a	Age	No. of embryos transferred	Results
1	Ham's F-10 +50% HFF	32	5	Triplets delivered
2	Ham's F-10 +50% HFF	43	1	Singleton delivered
3	Ham's F-10 +50% HPF	51	3	Clinical abortion
4	Ham's F-10 +50% HPF	33	5	Clinical abortion
5	TCM-199 +50% HPF	32	10	Clinical abortion
6	TCM-199 +50% HPF	31	5	Twins delivered
7	TCM-199 +50% HPF	29	3	Singleton delivered

^aHFF: human follicular fluid; HPF: human peritoneal fluid.

It seems that there is a relationship between follicular size in the human ovary and oocytes capable of resuming meiosis *in vitro* (Tsuji *et al.*, 1985). In the follicular phase, the percentage

of oocyte maturation in the large-follicle group (9–15 mm diameter; 34.5%) was significantly ($P < 0.05$) higher than that in the small-follicle group (3–4 mm; 8.8%). It has been reported that the unstimulated human oocyte appears to have a size-dependent ability to resume meiosis and complete maturation, indicating a significant difference in oocytes measuring 86–105 μm (22.2%) versus those measuring 106–125 μm (60.0%) (Durinzi *et al.*, 1995).

Table IV. Maturation and fertilization rates of human oocytes derived from unstimulated cycles^a

	Maturation (%) ^b (<i>n</i> = 199)	Fertilization (%) ^c (<i>n</i> = 27)
No. of oocytes	158 (79.4)	25 (92.6)

^aOocytes were cultured in TCM-199 + 20% fetal bovine serum supplemented with gonadotrophins (10 IU/ml pregnant mare's serum gonadotrophin and 10 IU/ml human chorionic gonadotrophin). Because of ethical problems, some egg donors did not want their eggs to be mixed with donor spermatozoa. Thus only a small proportion of mature oocytes were used in the fertilization attempt, the remainder going to cryopreservation experiments.

^bExtrusion of first polar body.

^cFormation of both pronuclei.

An earlier study indicated that 42% (52/124) of immature oocytes displayed GVBD and 31% (39/124) progressed to metaphase II when the oocytes were cultured with TCM-199 or Ham's F-10 supplemented with 10% fetal calf serum for 42 h (Shea *et al.*, 1975). The maturation and fertilization rates of immature human oocytes were 63% and 60%, respectively, when using human follicular fluid or peritoneal fluid as supplements in the maturation medium (Cha *et al.*, 1989). The first live birth from immature human oocytes obtained from unstimulated ovaries was in our ovum donation programme (Cha *et al.*, 1991). Five embryos were transferred to a woman with premature ovarian failure and she delivered healthy triplet girls. Shortly thereafter, from 21 transfer cycles, viable embryos were derived from follicular immature oocytes matured *in vitro* in Ham's F-10 medium supplemented with 50% follicular fluid or 50% peritoneal fluid, resulting in another two live births (Cha *et al.*, 1992). In our centre, the human ovum donation programme has subsequently obtained six more pregnancies by using follicular immature oocytes derived from donors who underwent tuboplasty or Caesarean section (Cha, 1995; Table III). Recently, we have been using TCM-199 + 20% fetal bovine serum (FBS) and supplemented with gonadotrophins [10 IU/ml pregnant mare's serum gonadotrophin (PMSG) and 10 IU/ml HCG], resulting in ~80% maturation rate and >90% fertilization rate when the

germinal vesicle oocytes were retrieved from unstimulated ovaries (Table IV).

The time courses of GVBD and maturation of the oocytes were different between the immature oocytes retrieved from stimulated and unstimulated ovaries (Figures 4 and 5). Most oocytes matured to metaphase II 45 h after culture *in vitro* when the immature germinal vesicle oocytes were retrieved from the patients undergoing Caesarean section (Figure 5). However, when the immature germinal vesicle oocytes were recovered from stimulated cycles, most oocytes reached metaphase II at 30 h after culture *in vitro*. The final percentages of the oocyte maturation were not different between the immature oocytes derived from stimulated (75.0% = 27/36) and unstimulated (77.5% = 31/40) cycles. However, the percentage of metaphase II oocytes was significantly higher for germinal vesicle oocytes derived from stimulated rather than unstimulated ovaries (Gómez *et al.*, 1993a). Combination of the results from Toth *et al.* (1994a,b) also indicated that the maturation rates are different between the immature human oocytes retrieved from stimulated and unstimulated cycles. These differences might be due to the use of different maturation medium and different culture supplements during maturation in each research group. Ninety per cent (18/20) of germinal vesicle oocytes retrieved from patients undergoing Caesarean section presented a nucleolus in the oocyte germinal vesicle, but only 20% (2/10) of oocytes retrieved from stimulated cycles contained a nucleolus in the germinal vesicle. The profile of protein synthesis is also different in these two groups (Chian *et al.*, 1997). These results suggest that the differences between the germinal vesicle oocytes retrieved from stimulated and unstimulated cycles may subsequently relate to time courses of GVBD and maturation of oocytes cultured *in vitro* in these two groups.

The disadvantages of ovulation induction include the side-effects from the gonadotrophins, which include weight gain, bloating, nausea, vomiting, mood swings, ovarian hyperstimulation with possible hospitalization, and the unknown long-term cancer risks (Bergh and Navot, 1992). Therefore, many attempts have been made to perform IVF without the use of exogenous gonadotrophins (natural cycle) to induce multiple follicles. One very attractive possibility for enhancement of the success of natural cycle IVF is the combination of oocyte maturation techniques with immature oocyte retrieval, and this technique has yielded live births (Paulson *et al.*, 1994; Russell *et al.*, 1997). Recently it has been reported that an average of 11 oocytes per patient (from unstimulated ovaries) were retrieved transvaginally with ultrasound guidance, which suggests that unstimulated immature oocyte retrieval can be performed successfully in patients




Figure 5. The time course of human oocyte maturation [polar body (PB) exclusion] during culture *in vitro*. There was no significant difference in maturation rates between the oocytes retrieved from stimulated (—●—) and unstimulated (---○---) ovaries.

who are candidates with different diagnoses for assisted reproductive treatment (Russell *et al.*, 1997). This indicates that the immature oocyte collection and in-vitro maturation procedures could possibly replace the standard stimulated IVF cycle in the near future.

Oocytes from polycystic ovarian syndrome patients

Patients with polycystic ovarian syndrome (PCOS) are characterized by abnormal endocrine parameters, anovulation, numerous antral follicles within their ovaries and frequently infertility (Goldzieher and Green, 1962). PCOS patients may be extremely sensitive to exogenous gonadotrophin and may be at risk of ovarian hyperstimulation syndrome when treated with gonadotrophins for assisted reproduction. Immature oocyte recovery could be developed as a new method for the treatment of patients with infertility due to PCOS because the oocytes of these patients retain their maturational and developmental competence (Trounson *et al.*, 1994). They reported that 65% of oocytes cultured in medium with gonadotrophins, oestrogen and fetal calf serum matured to metaphase II by 43–47 h, and 81% were matured at 48–54 h of culture. Of the inseminated oocytes, 34% fertilized and 56% of the cultured pronuclear oocytes cleaved to eight cells or more. A pregnancy and the first birth of a normal baby occurred in one of the anovulatory PCOS patients receiving an abbreviated steroid replacement protocol after embryo transfer. A second birth was achieved by in-vitro maturation of primary oocytes derived from a PCOS patient, combined with ICSI and assisted hatching (Barnes *et al.*, 1995), indicating that the developmental capacity of

primary oocytes is higher in regular cycling women than in irregular cycling and anovulatory women with PCOS (Barnes *et al.*, 1996).

In our centre, the PCOS patients are diagnosed when the following conditions are present: physical characteristics such as obesity and hirsutism, menstrual disturbances (oligomenorrhoea, amenorrhoea), LH/FSH ratio >3.0 and the characteristic polycystic ovarian appearance at ultrasonographic assessment. For maturation *in vitro* of immature oocytes retrieved from these PCOS patients, TCM-199 + 20% FBS and supplemented with 10 IU/ml PMSG and 10 IU/ml HCG were used. It has been found that the maturation rate of immature oocytes retrieved from PCOS patients is significantly lower than those recovered from normal menstrual cycle patients (K.Y.Cha *et al.*, unpublished data). The results indicate that 60.0% (499/832) of oocytes cultured *in vitro* matured to metaphase II and that 73.0% (364/499) of the oocytes fertilized by ICSI and 89.9% (187/208) of the cultured pronuclear oocytes cleaved to 2- and 4-cell stage (Cha *et al.*, 1996). Up to now, a total of 16 pregnancies have been achieved from PCOS patients in our centre (Table V). These indicate that immature oocyte recovery from women with infertility due to PCOS can be used with this therapy for assisted reproduction.

Cryopreservation of immature oocytes

Successful cryopreservation of the immature oocytes would widen the possibility of establishing an ovum bank. Oocyte cryopreservation has the potential to overcome many of the legal and ethical problems associated with embryo cryopreservation and provides more options for patient treatment. A clinical application of cryopreservation of immature oocytes would be to store oocytes from patients who risk the loss of ovarian function. Oocytes are very large cells with well-defined structures which do not withstand temperature alterations without impairment of their fertilizing and developmental capacity. During freezing and thawing, the oocyte changes with rearrangement of the cytoskeleton and of the meiotic spindle, clumping and migration of the cortical granules

and hardening of the zona pellucida (Vincent and Johnson, 1992; George *et al.*, 1994; Van Blerkom and Davis, 1994). Although successful pregnancies after cryopreservation of metaphase II oocytes have been reported (Chen, 1986; Al-Hasani *et al.*, 1987; Van Uem *et al.*, 1987), the success rate is still low. It has been suggested that the microtubular spindle of metaphase II oocytes is sensitive to temperature changes (Van der Elst *et al.*, 1988). Thus, chromatid non-disjunction may occur during cooling and it would result in aneuploidy after fertilization (Pickering *et al.*, 1990). Freezing of immature oocytes may solve these problems. The advantage of freezing at prophase I is that the procedure can be achieved before spindle formation in the oocytes.

Production of live young after immature oocyte freezing has been successful in mice (Candy *et al.*, 1994). Human germinal vesicle oocytes can survive and mature *in vitro* after freeze-thawing (Mandelbaum *et al.*, 1988; Toth *et al.*, 1994a,b; Son *et al.*, 1996; Park *et al.*, 1997): 55.1% (54/98) of oocytes survived and 59.3% (32/54) of oocytes matured to metaphase II following freeze-thawing, 42.9% (6/14) of oocytes fertilized and 16.7% (1/6) of fertilized oocytes cleaved (Table VI), suggesting that immature human oocytes are capable of meiotic maturation, fertilization and development following cryopreservation (Son *et al.*, 1996; Park *et al.*, 1997). However, the freeze-thawing procedures have detrimental effects on the maturation of human oocytes and subsequent early developmental capacity. It has been indicated that human oocytes matured *in vitro* after cryopreservation at the germinal vesicle stage also showed an increased incidence of chromosomal and spindle abnormalities, suggesting that these abnormalities may impair the capacity for further development of embryos derived from frozen-thawed oocytes (Park *et al.*, 1997). Furthermore, when in-vitro matured germinal vesicle oocytes after cryopreservation were inseminated by either conventional IVF or ICSI, the fertilization rate was higher in the ICSI group (64.3%) than the conventional IVF group (27.3%) (Park *et al.*, 1996). This result suggests that the reduced rate of fertilization after freeze-thawing may be due to hardening of the zona pellucida.

Table V. Pregnancy outcome of immature oocytes derived from PCOS patients^a

No. of cases	No. of oocytes retrieved	No. of oocytes cultured	No. of oocytes matured (%)	No. of oocytes fertilized (%) ^b	No. of embryos transferred (%) ^c	Pregnancies (%) ^d
72	910	832	499 (60.0)	400 (80.2)	306 (76.5)	16 (25.0)

^aData modified from Cha *et al.* (1996).

^bAbnormally fertilized oocytes ($n = 36$) were included.

^cNo. of embryos transferred/no. of fertilized oocytes cultured = 187/208. Mean no. of embryos transferred was 4.7 per patient.

^dNo. of pregnant patients/no. of patients who had embryos transferred = 16/64; one of these pregnancies was a twin pregnancy, whilst the rest were singletons.

Table VI. Comparison of maturation, fertilization and cleavage rates between cryopreserved immature human oocytes and control^a

	Control	Frozen-thawed at germinal vesicle stage ^b
No. of oocytes	82	98
No. surviving (%)	82 (100)	54 (55.1)
No. undergoing maturation (%)	63 (76.8)	32 (59.3)
No. undergoing insemination	21	14
No. fertilized (%)	19 (90.5)	6 (42.9)
No. cleaved (%)	18 (94.7)	1 (16.7)

^aData modified from Son *et al.* (1996).^bSignificantly different from control ($P < 0.01$).

Early studies indicated that restoration of fertility could be obtained by reimplantation of frozen-thawed ovarian tissue (Parrott, 1960) and that only primordial follicles survived and allowed in-vitro development following the cryopreservation process (Deanesly, 1954; Green *et al.*, 1956; Parkes, 1958). Since these studies, cryopreservation techniques have been much improved. Recently the technique of reimplantation of frozen-thawed ovarian tissue has been applied to several species resulting in live offspring (Carroll and Gosden, 1993; Gosden *et al.*, 1994). On the other hand, it has been known that immature oocytes from early preantral follicles from prepubertal mice matured, fertilized, and the embryos developed *in vitro* in a simplified culture system (Cortvindt *et al.*, 1996a). They also indicated that cryopreserved whole early preantral follicles matured *in vitro* and responded to stimulation with HCG as the control group and that these cryopreserved, in-vitro matured oocytes had the potential to fertilize and develop to hatched blastocysts (Cortvindt *et al.*, 1996b). These experiments suggest that freezing of whole early follicles may circumvent potential injuries to chromosomal material, cytoplasmic constituents and to the sperm penetration potential of the zona and oolemma. Cryopreserving isolated whole follicles and maturing the oocytes in the follicles *in vitro* may provide more options for patient treatment.

Conclusions

The cytoplasm of the oocyte is of key interest in oocyte maturation. The best way to improve embryo quality is to improve oocyte quality. Addition of gonadotrophins (FSH, LH) to the maturation medium is beneficial to cytoplasmic maturation of immature human oocytes. Oocytes derived from stimulated and unstimulated ovaries can be matured and fertilized *in vitro* even though the fertilization rates are different. Immature oocytes retrieved from PCOS patients also can be matured and fertilized *in vitro*. This indicates that immature oocytes recovered from women with infertility due to PCOS can be used with this therapy for

assisted reproduction. Immature germinal vesicle oocytes are also capable of meiotic maturation, fertilization and development *in vitro* following cryopreservation.

Acknowledgements

We are grateful to Dr Hugh J. Clarke, Department of Obstetrics and Gynecology, McGill University, Canada, for his helpful comments on early drafts of the manuscript. We gratefully acknowledge Mr Simon Phillips, McGill Reproductive Center, for his assistance in the preparation of this manuscript.

References

- Acharya, U., Small, J. and Randall, J. *et al.* (1992) Prospective study of short and long regimens of gonadotropin-releasing hormone agonist in *in vitro* fertilization program. *Fertil. Steril.*, **57**, 815–818.
- Al-Hasani, S., Diedrich, K. and Van der Ven, H. *et al.* (1987) Cryopreservation of human oocyte. *Hum. Reprod.*, **2**, 695–700.
- Alsallil, M., Thornton, S. and Fleming, S. (1997) The effect of the anaesthetic, Propofol, on *in vitro* oocyte maturation, fertilization and cleavage in mice. *Hum. Reprod.*, **12**, 1271–1274.
- Artley, J.K., Braude, P.R. and Johnson, M.H. (1992) Gene activity and cleavage arrest in human pre-embryos. *Hum. Reprod.*, **7**, 1014–1021.
- Ashwood-Smith, M. and Edwards, R.G. (1996) DNA repair by oocytes. *Mol. Hum. Reprod.*, **2**, 46–51.
- Bachvarova, R. (1974) Incorporation of tritiated adenosine into mouse ovum RNA. *Dev. Biol.*, **40**, 52–58.
- Bachvarova, R. and DeLeon, V. (1980) Polyadenylated RNA of mouse ova and loss of maternal RNA in early development. *Dev. Biol.*, **74**, 1–8.
- Baker, T.G. (1963) Quantitative and cytological study of germ cells in human ovaries. *Proc. R. Soc.*, **B158**, 417–433.
- Barnes, F.L., Crombie, A. and Gardner, D.K. *et al.* (1995) Blastocyst development and birth after *in vitro* maturation of human primary oocytes, intracytoplasmic sperm injection and assisted hatching. *Hum. Reprod.*, **10**, 3243–3247.
- Barnes, F.L., Kausche, A. and Tiglias, J. *et al.* (1996) Production of embryos from *in vitro*-matured primary human oocytes. *Fertil. Steril.*, **65**, 1151–1156.
- Barnett, D. and Bavister, B. (1996) What is the relationship between the metabolism of preimplantation embryos and their developmental competence? *Mol. Reprod. Dev.*, **43**, 105–133.
- Battaglia, D., Goodwin, P. and Klein, N. *et al.* (1996a) Influence of maternal age on meiotic spindle assembly in oocytes from naturally cycling women. *Hum. Reprod.*, **11**, 2217–2222.
- Battaglia, D.E., Klein, N.A. and Soules, M.R. (1996b) Changes in centrosomal domains during meiotic maturation in the human oocyte. *Mol. Hum. Reprod.*, **2**, 845–851.
- Bergh, P.A. and Navot, D. (1992) Ovarian hyperstimulation syndrome: a review of pathophysiology. *J. Assist. Reprod. Genet.*, **9**, 429–438.
- Berridge, M.J. (1996) Regulation of calcium spiking in mammalian oocytes through a combination of inositol triphosphate-dependent entry and release. *Mol. Hum. Reprod.*, **2**, 386–388.
- Berridge, M.J. and Irvine, R.F. (1984) Inositol triphosphate, a novel second messenger in cellular signal transduction. *Nature*, **312**, 315–321.
- Bezprozvanny, I., Watras, J. and Ehrlich, B.E. (1991) Bell shaped calcium-response curves of Ins(1,4,5)P₃- and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature*, **351**, 751–754.
- Bloom, A.M. and Mukherjee, B.B. (1972) RNA synthesis in maturing mouse oocytes. *Expl. Cell Res.*, **74**, 577–582.
- Bomse-Helmreich, O., Huyen, L.V.N. and Durand-Gassel, I. *et al.* (1987) Mature and immature oocytes in large and medium follicles after clomiphene citrate and human menopausal gonadotropin stimulation without human chorionic gonadotropin. *Fertil. Steril.*, **48**, 596–604.
- Bongso, A., Chye, N.S. and Ratnam, S. *et al.* (1988) Chromosome anomalies in human oocytes failing to fertilize after insemination *in vitro*. *Hum. Reprod.*, **3**, 645–649.

- Boué, J.G. and Boué, A. (1973) Increased frequency of chromosomal abnormalities in abortions after induced ovulation. *Lancet*, **1**, 679–680.
- Bourguignon, L.Y.W., Jin, H. and Iida, N. *et al.* (1993) The involvement of ankyrin in the regulation of inositol 1,4,5-trisphosphate receptor-mediated internal Ca^{2+} release from C storage vesicles in mouse T-lymphoma cells. *J. Biol. Chem.*, **268**, 7290–7297.
- Braude, P., Bolton, V. and Moore, S. (1988) Human gene expression first occurs between the four- and eight-cell stage of preimplantation development. *Nature*, **332**, 459–461.
- Braude, P.R., Pickering, S.J. and Winston, N.J. *et al.* (1990) The development of the human preimplantation embryo. *I Proceedings III Renier de Graaff Symposium. From Ovulation to Implantation*. Elsevier, Amsterdam, pp. 251–261.
- Braw-Tal, R. (1994) Expression of mRNA for follistatin and inhibin/activin subunits during follicular growth and atresia. *J. Mol. Endocrinol.*, **13**, 253–264.
- Brillantes, A.M.B., Ondrias, K. and Scott, A. *et al.* (1994) Stabilization of calcium-release channel (ryanodine receptor) function by FK506 binding protein. *Cell*, **77**, 513–523.
- Cameron, A.M., Steiner, J.P. and Sabatini, D.M. *et al.* (1995) Immunophilin FK506 binding protein associate with inositol 1,4,5-trisphosphate receptor mediates calcium influx. *Proc. Natl. Acad. Sci. USA*, **92**, 1784–1788.
- Candy, C.J., Wood, M.J. and Whittingham, D.G. *et al.* (1994) Cryopreservation of immature mouse oocytes. *Hum. Reprod.*, **9**, 1738–1742.
- Canipari, R., Palombi, F. and Riminucci, M. *et al.* (1984) Early programming of maturation competence in mouse oogenesis. *Dev. Biol.*, **102**, 519–524.
- Carroll, J. and Gosden, R.G. (1993) Transplantation of frozen-thawed mouse primordial follicles. *Hum. Reprod.*, **8**, 1163–1167.
- Carroll, J., Jones, K.T. and Whittingham, D.G. (1996) Ca^{2+} release and the development of Ca^{2+} release mechanisms during oocyte maturation: a prelude to fertilization. *Rev. Reprod.*, **1**, 137–143.
- Cha, K.Y. (1992) In vitro fertilization using immature follicular oocytes harvested from ovarian tissue. In Behrman, S.J., Patton, G.W. Jr and Holtz, G. (eds), *Progress in Infertility*, **7**, 99–112.
- Cha, K.Y. (1995) Clinical use of immature oocytes. *28th Annual Postgraduate Program. American Society of Reproductive Medicine*, pp. 97–114.
- Cha, K.Y., Yoon, S.H. and Ko, J.J. *et al.* (1989) Pregnancy after *in vitro* fertilization of human follicular oocytes collected from non-stimulated cycle, their culture *in vitro* and their transfer according to the donor oocyte program. *Proceedings of Annual Meeting of American Fertility Society*, Abstr. O-001.
- Cha, K.Y., Koo, J.J. and Ko, J.J. *et al.* (1991) Pregnancy after *in vitro* fertilization of human follicular oocytes collected from nonstimulated cycles, their culture *in vitro* and their transfer in a donor oocyte program. *Fertil. Steril.*, **55**, 109–113.
- Cha, K.Y., Do, B.R. and Chi, H.J. *et al.* (1992) Viability of human follicular oocytes collected from unstimulated ovaries and matured and fertilized *in vitro*. *Reprod. Fertil. Dev.*, **4**, 695–701.
- Cha, K. Y., Chung, H. M. and Han, S. Y. *et al.* (1996) Successful *in vitro* maturation, fertilization and pregnancy by using immature follicular oocytes collected from unstimulated polycystic ovarian syndrome patients. *Proceedings of Annual Meeting of American Society for Reproductive Medicine*, Abstr. O-044.
- Chapman, D. and Wolgemuth, D. (1992) Identification of a mouse B-type cyclin which exhibits developmentally regulated expression in the germ line. *Mol. Reprod. Dev.*, **33**, 259–269.
- Chapman, D. and Wolgemuth, D. (1993) Isolation of the murine cyclin B2 cDNA and characterization of expression of the B1 and B2 cyclins during oogenesis, spermatogenesis and early embryogenesis. *Development*, **118**, 229–240.
- Chen, C. (1986) Pregnancy after human oocyte cryopreservation. *Lancet*, **1**, 884–886.
- Chian, R.C. and Sirard, M.A. (1995) Effects of cumulus cells and follicle-stimulating hormone during *in vitro* maturation on parthenogenetic activation of bovine oocytes. *Mol. Reprod. Dev.*, **42**, 425–431.
- Chian, R.C., Niwa, K. and Nakahara, H. (1992) Effect of sperm penetration *in vitro* on completion of first meiosis of bovine oocytes arrested at various stages in culture. *J. Reprod. Fertil.*, **96**, 73–78.
- Chian, R.C., Niwa, K. and Sirard, M. A. (1994) Effects of cumulus cells on male pronuclear formation and subsequent early development of bovine oocytes *in vitro*. *Theriogenology*, **41**, 1499–1508.
- Chian, R.C., Park, S.E. and Park, E.H. *et al.* (1997) Molecular and structural characteristics between immature human oocytes retrieved from stimulated and unstimulated ovaries. *Proceedings of 10th World Congress of IVF and AR*, Monduzzi, Editore, Bologna, pp. 315–319.
- Chiba, K., Kado, R.T. and Jaffe, L.A. (1990) Development of calcium release mechanism during starfish oocyte maturation. *Dev. Biol.*, **140**, 300–306.
- Cioffi, J.A., Van Blerkom, J. and Antczak, M. *et al.* (1997) The expression of leptin and its receptors in pre-ovulatory human follicles. *Mol. Hum. Reprod.*, **3**, 467–472.
- Clarke, H.J. and Masui, Y. (1983) The induction of reversible and irreversible chromosome decondensation by protein synthesis inhibition during meiotic maturation of mouse oocytes. *Dev. Biol.*, **97**, 291–301.
- Coetsier, T., Dhont, M. and De Sutter, P. *et al.* (1992) Propofol anaesthesia for ultrasound guided oocyte retrieval: accumulation of the anaesthetic agent in follicular fluid. *Hum. Reprod.*, **7**, 1422–1424.
- Cortvrindt, R., Smits, J. and Van Steirteghem, A. (1996a) In-vitro maturation, fertilization and embryo development of immature oocytes from early preantral follicles from prepubertal mice in a simplified culture system. *Hum. Reprod.*, **11**, 2656–2666.
- Cortvrindt, R., Smits, J. and Van Steirteghem, A. (1996b) A morphological and functional study of the effect of slow freezing followed by complete in-vitro maturation of primary mouse ovarian follicles. *Hum. Reprod.*, **11**, 2648–2655.
- Crosby, I.M., Gandolfi, F. and Moor, R.M. (1988) Control of protein synthesis during early cleavage of sheep embryos. *J. Reprod. Fertil.*, **82**, 769–775.
- Cross, P. C. and Brinster, R. L. (1970) In vitro development of mouse oocytes. *Biol. Reprod.*, **3**, 298–307.
- Dandekar, P.V., Martin, M.C. and Glass, R.H. (1991) Maturation of immature oocytes by coculture with granulosa cells. *Fertil. Steril.*, **55**, 95–99.
- Davis, D. (1985) Culture and storage of pig embryos. *J. Reprod. Fertil.*, **38** (Suppl.), 115–124.
- Deanesly, R. (1954) Immature rat ovaries grafted after freezing and thawing. *J. Endocrinol.*, **11**, 197–200.
- Dekel, N. (1988) Spatial relationship of follicular cells in control of meiosis. In Haseltine, F.P. and First, N.L. (eds), *Meiotic Inhibition, Molecular Control of Meiosis: Progress in Clinical and Biological Research*. Alan, R. Liss, New York, pp. 87–101.
- Dekel, N., Lawrence, T. S., Gilula, N. B. and Beers, W. H. (1981) Modulation of cell-to-cell communication in the cumulus-oocyte complex and regulation of oocyte maturation by LH. *Dev. Biol.*, **86**, 352–362.
- Delhanty, J. and Handyside, A. (1995) The origin of genetic defects in the human and their detection in the preimplantation embryo. *Hum. Reprod. Update*, **1**, 201–215.
- Depypere, H.T., Dhont, M. and De Sutter, P. *et al.* (1991) The influence of propofol on *in vitro* fertilization in mice. *Hum. Reprod.*, **6** (Suppl. 1), 151 [Abstr. No. 127].
- De Sutter, P., Dhont, M. and Verschraegen-Spae, M.R. *et al.* (1991a) Chromosome analysis in human oocytes non fertilized *in vitro*: a mathematical model for the estimation of the first meiotic non-disjunction frequency. *Hum. Reprod.*, **6**, 550–554.
- De Sutter, P., Dhont, M. and Vanluchene, E. *et al.* (1991b) Correlations between follicular fluid steroid analysis, maturity and cytogenetic analysis of human oocytes which remained unfertilized after *in vitro* fertilization. *Fertil. Steril.*, **55**, 958–963.
- De Sutter, P., Dhont, M. and Vandekerckhove, D. (1992) Hormonal stimulation for *in vitro* fertilization: a comparison of fertilization rates and cytogenetic findings in unfertilized oocytes. *J. Assist. Reprod. Genet.*, **9**, 254–258.
- Dimfeld, M., Goldman, S. and Gonen, Y. *et al.* (1993) Functional differentiation in progesterone secretion by granulosa versus cumulus cells in the human preovulatory follicle and the effect of different induction of ovulation protocols. *Fertil. Steril.*, **60**, 1025–1030.
- Djalali, M., Rosenbusch, B. and Wolf, M. *et al.* (1988) Cytogenetics of unfertilized human oocytes. *J. Reprod. Fertil.*, **84**, 647–652.
- Downs, S.M. (1995) The influence of glucose, cumulus cells, and metabolic coupling on ATP levels and meiotic control in the isolated mouse oocytes. *Dev. Biol.*, **167**, 502–512.
- Downs, S.D., Coleman, D.L. and Ward-Bailey, P.F. *et al.* (1985) Hypoxanthine is the principal inhibitor of murine oocyte maturation in a low molecular weight fraction of porcine follicular fluid. *Proc. Natl. Acad. Sci. USA*, **82**, 454–458.

- Downs, S.M., Daniel, S.A.J. and Eppig, J.J. (1988) Induction of maturation in cumulus cell-enclosed mouse oocytes by follicle-stimulating hormone and epidermal growth factor: evidence for a positive stimulus of somatic cell origin. *J. Exp. Zool.*, **245**, 86–96.
- Dunphy, W., Brizuela, L. and Beach, D. *et al.* (1988) The *Xenopus cdc2* protein is a component of MPF, a cytoplasmic regulator of mitosis. *Cell*, **54**, 423–431.
- Durinzi, K.L., Saniga, E.M. and Lanzendorf, S.E. (1995) The relationship between size and maturation *in vitro* in the unstimulated human oocyte. *Fertil. Steril.*, **63**, 404–406.
- Durinzi, K.L., Wentz, A.C. and Saniga, E.M. *et al.* (1997) Follicle stimulating hormone effects on immature human oocytes: *in vitro* maturation and hormone production. *J. Assist. Reprod. Genet.*, **14**, 199–204.
- Edirisinghe, W., Junk, S.M. and Matson, P.L. *et al.* (1997) Birth from cryopreserved embryos following *in-vitro* maturation of oocytes and intracytoplasmic sperm injection. *Hum. Reprod.*, **12**, 1056–1058.
- Edwards, R.G. (1965a) Maturation *in vitro* of mouse, sheep, cow, pig, rhesus monkey and human ovarian oocytes. *Nature*, **208**, 349–351.
- Edwards, R.G. (1965b) Maturation *in vitro* of human ovarian oocytes. *Lancet*, **ii**, 926–929.
- Edwards, R.G. and Beard, H.K. (1997) Oocyte polarity and cell determination in early mammalian embryos. *Mol. Hum. Reprod.*, **3**, 863–905.
- Edwards, R.G., Bavister, B.D. and Steptoe, P.C. (1969) Early stages of fertilization *in vitro* of human oocytes matured *in vitro*. *Nature*, **221**, 632–635.
- Eppig, J.J. (1993) Regulation of mammalian oocyte maturation. In Adashi, E.Y. and Leung, P.C.K. (eds), *The Ovary*. Raven Press, New York, pp. 185–208.
- Eppig, J.J., Freter, R. R. and Ward-Bailey, P. F. *et al.* (1983) Inhibition of oocyte maturation in the mouse: participation of cAMP, steroids, and a putative maturation inhibitory factor. *Dev. Biol.*, **100**, 39–49.
- Eppig, J.J., Ward-Bailey, P.F. and Coleman, D.L. (1985) Hypoxanthine and adenosine in murine ovarian follicular fluid: concentrations and activity in maintaining oocyte meiotic arrest. *Biol. Reprod.*, **33**, 1041–1049.
- Eppig, J.J., Schroeder, A.C. and O'Brien, M.J. (1992) Developmental capacity of mouse oocytes matured *in vitro*: effect of gonadotrophic stimulation, follicular origin and oocyte size. *J. Reprod. Fertil.*, **95**, 119–127.
- Evans, T., Rosenthal, E.T. and Youngblom, J. *et al.* (1983) Cyclin: a protein specified by maternal mRNA in sea urchin eggs that is destroyed at each cleavage division. *Cell*, **33**, 389–396.
- Fagbohun, C.F. and Downs, S.M. (1991) Metabolic coupling and ligand-stimulated meiotic maturation in the mouse oocyte-cumulus cell complex. *Biol. Reprod.*, **45**, 851–859.
- Ferrell, J.E., Wu, M. and Gerhardt, J.C. *et al.* (1991) Cell cycle tyrosine phosphorylation of p34^{cdc2} and MAP-kinase homolog in *Xenopus* oocytes and eggs. *Mol. Cell Biol.*, **11**, 1965–1971.
- Flach, G., Johnson, M. and Braude, P. *et al.* (1982) The transition from maternal to embryonic control in the 2-cell mouse embryo. *EMBO J.*, **1**, 681–686.
- Frei, R., Schultz, G. and Church, R. (1989) Qualitative and quantitative changes in protein synthesis occur at the 8–16 cell stage of embryogenesis in the cow. *J. Reprod. Fertil.*, **86**, 637–641.
- Frydman, R., Belaisch-Allart, J. and Parneix, I. *et al.* (1988) Comparison between flare up and down regulation effects of luteinizing hormone-releasing hormone agonists in an *in vitro* fertilization program. *Fertil. Steril.*, **50**, 471–475.
- Fujiwara, T., Nakade, K. and Shirakawa, H. *et al.* (1993) Development of inositol triphosphate-induced calcium release mechanisms during maturation of hamster oocytes. *Dev. Biol.*, **156**, 69–79.
- Fukui, Y. (1990) Effect of follicle cells on the acrosome reaction, fertilization, and developmental competence of bovine oocytes matured *in vitro*. *Mol. Reprod. Dev.*, **26**, 40–46.
- Fulka Jr, J., Kalab, P. and First, N.L. *et al.* (1997) Damaged chromatin does not prevent the exit from metaphase I in fused mouse oocytes. *Hum. Reprod.*, **12**, 2473–2476.
- Furuichi, T., Yoshikawa, S. and Miyawaki, A. *et al.* (1989) Primary structure and functional expression of the inositol 1,4,5-trisphosphate-binding protein P₄₀₀. *Nature*, **342**, 32–38.
- Gandolfi, F. and Moor, R. (1987) Stimulation of early embryonic development in the sheep by co-culture with oviduct epithelial cells. *J. Reprod. Fertil.*, **81**, 23–28.
- Garcia, J.E., Padilla, S.L. and Bayati, J. *et al.* (1990) Follicular phase gonadotropins: a better alternative for ovulation induction in *in vitro* fertilization. *Fertil. Steril.*, **53**, 302–305.
- Gardner, R.L. (1996) Can developmentally significant spatial patterning of the egg be discounted in mammals? *Hum. Reprod. Update*, **2**, 3–27.
- Gautier, J., Norbury, C. and Lohka, M. *et al.* (1988) Purified maturation promoting factor contains the product of a *Xenopus* homolog of the fission yeast cell cycle control gene *cdc2*. *Cell*, **54**, 433–439.
- Gautier, J., Minshull, J. and Lohka, M. *et al.* (1990) Cyclin is a component of maturation promoting factor from *Xenopus*. *Cell*, **60**, 487–494.
- George, M.A., Johnson, M.H. and Howlett, S.K. (1994) Assessment of the developmental potential of frozen-thawed mouse oocytes. *Hum. Reprod.*, **9**, 130–136.
- Gerhart, J., Wu, M. and Kirschner, M. (1984) Cell cycle dynamics of an M-phase specific cytoplasmic factor in *Xenopus laevis* oocytes and eggs. *J. Cell Biol.*, **98**, 1247–1255.
- Gifford, D.J., Fleetham, J.A. and Mahadevan, M.M. *et al.* (1987) Protein synthesis in human unfertilized eggs. *Gamete Res.*, **18**, 97–107.
- Goldman, D., Kiessling, A. and Millette, C. *et al.* (1987) Expression of c-mos RNA in germ cells of male and female mice. *Proc. Natl. Acad. Sci. USA*, **84**, 4509–4513.
- Goldman, S., Dirnfeld, M. and Gonen, Y. *et al.* (1993) Different morphology and proliferative ability of cumulus and granulosa cells originating from cystic follicles aspirated from stimulated *in vitro* fertilization patients. *Fertil. Steril.*, **59**, 601–605.
- Goldzieher, J.W. and Green, J.A. (1962) The polycystic ovary. I. Clinical and histologic features. *J. Clin. Endocrinol. Metab.*, **22**, 325–338.
- Gómez, E., Tarín, J. J. and Pellicer, A. (1993a) Oocyte maturation in humans: the role of gonadotropins and growth factors. *Fertil. Steril.*, **60**, 40–46.
- Gómez, E., de los Santos, M.J. and Ruiz, A. *et al.* (1993b) Effects of epidermal growth factor in the final stages of nuclear and cytoplasmic oocyte maturation in humans. *Hum. Reprod.*, **8**, 691–694.
- Gosden, R.G., Baird, D.T. and Wade, J.C. *et al.* (1994) Restoration of fertility to oophorectomized sheep by ovarian autografts stored at –196°C. *Hum. Reprod.*, **9**, 597–603.
- Gras, L., McBain, J. and Trounson, A. *et al.* (1992) The incidence of chromosomal aneuploidy in stimulated and unstimulated (natural) uniseminated human oocytes. *Hum. Reprod.*, **7**, 1396–1401.
- Green, S.H., Smith, A.U. and Zuckerman, S. (1956) The numbers of oocytes in autografts after freezing and thawing. *J. Endocrinol.*, **13**, 330–334.
- Greenblatt, E.M., Meriano, J.S. and Casper, R.F. (1995) Type of stimulation protocol affects oocyte maturity, fertilization rate, and cleavage rate after intracytoplasmic sperm injection. *Fertil. Steril.*, **64**, 557–563.
- Gregory, L. and Leese, H. J. (1996) Determinants of oocyte and preimplantation embryo quality: metabolic requirements and the potential role of cumulus cells. *Hum. Reprod.*, **11**, 96–102.
- Hashimoto, N. and Kishimoto, T. (1988) Regulation of meiotic metaphase by a cytoplasmic maturation-promoting factor during mouse oocyte maturation. *Dev. Biol.*, **126**, 242–252.
- Heikinheimo, O., Lanzendorf, S.E. and Baka, S.G. *et al.* (1995) Cell cycle genes c-mos and cyclin-B1 are expressed in a specific pattern in human oocytes and preimplantation embryos. *Mol. Hum. Reprod.*, **10**, 699–707.
- Heikinheimo, O., Toner, J.P. and Lanzendorf, S.E. *et al.* (1996) Messenger ribonucleic acid kinetics in human oocytes-effects of *in vitro* culture and nuclear maturational status. *Fertil. Steril.*, **65**, 1003–1008.
- Heller, D.T., Cahill, D.M. and Schultz, R.M. (1981) Biochemical studies of mammalian oogenesis: Metabolic cooperativity between granulosa cells and growing mouse oocytes. *Dev. Biol.*, **84**, 455–464.
- Herbert, M., Gillespie, J.I. and Murdoch, A.P. (1997) Development of calcium signalling mechanisms during maturation of human oocytes. *Mol. Hum. Reprod.*, **11**, 965–973.
- Hillensjo, T., Brannstrom, M. and Chare, S. *et al.* (1985) Oocyte maturation as regulated by follicular factors. *Ann. NY Acad. Sci.*, **442**, 73–79.
- Ho, Y., Doherty, A. and Schultz, R. (1994) Mouse preimplantation embryo development *in vitro*: Effects of sodium concentration in culture media on RNA synthesis and accumulation and gene expression. *Mol. Reprod. Dev.*, **38**, 131–141.
- Homburg, R., Armar, N.A. and Eshel, A. *et al.* (1988) Influence of serum luteinising hormone concentrations on ovulation, conception and early

- pregnancy loss in polycystic ovary syndrome. *Br. Med. J.*, **297**, 1024–1026.
- Howles, C.M., Macnamee, M.C. and Edwards, R.G. *et al.* (1986) Effect of high tonic levels of luteinizing hormone on outcome of *in vitro* fertilization. *Lancet*, **2**, 521–522.
- Hughes, E.G., Fedorkow, D.M. and Daya, S. *et al.* (1992) The routine use of gonadotropin-releasing hormone agonists prior to *in vitro* fertilization and gamete intrafallopian transfer: a meta-analysis of randomized controlled trials. *Fertil. Steril.*, **58**, 888–896.
- Hunter, T. (1991) Cell cycle gets more cyclins. *Nature*, **350**, 462–463.
- Iino, M. and Endo, M. (1992) Calcium-dependent immediate feedback control of inositol 1,4,5-trisphosphate-induced Ca^{2+} release. *Nature*, **360**, 76–78.
- Imthurn, B., Macas, E. and Rosselli, M. *et al.* (1997) Nuclear maturity and oocyte morphology after stimulation with highly purified follicle stimulating hormone compared to human menopausal gonadotrophin. *Hum. Reprod.*, **11**, 2387–2391.
- Iwamatsu, T. and Yanagimachi, R. (1975) Maturation *in vitro* of ovarian oocytes of prepubertal and adult hamsters. *J. Reprod. Fertil.*, **45**, 83–90.
- Jahn, C.L., Baran, M.M. and Bachvarova, R. (1976) Stability of RNA synthesized by the mouse oocyte during its major growth phase. *J. Exp. Zool.*, **197**, 161–172.
- Janssenswillen, C., Nagy, Z. and Steirteghem, A. (1995) Maturation of cumulus-free germinal vesicle-stage oocytes to metaphase II by coculture with monolayer Vero cells. *Hum. Reprod.*, **10**, 375–378.
- Janssenswillen, C., Christiaens, F. and Camu, F. *et al.* (1996) The effect of propofol on parthenogenetic activation, *in vitro* fertilization and early embryo development of mouse oocytes. *Hum. Reprod.* **12** (Nat. Suppl. 1), Abst. No. 38.
- Jaroudi, K.A., Hollanders, J.M.G. and Sieck, U.V. *et al.* (1997) Pregnancy after transfer of embryos which were generated from *in-vitro* matured oocytes. *Hum. Reprod.*, **12**, 857–859.
- Ji, Y.Z., Bomsel, M. and Jouannet, P. *et al.* (1996) Modifications of the human oocyte plasma membrane protein pattern during preovulatory maturation. *Mol. Reprod. Dev.*, **47**, 120–126.
- Jinno, M., Sadow, B.A. and Hodgen, G.D. (1989) Enhancement of the developmental potential of mouse oocytes matured *in vitro* by gonadotropins and ethylenediaminetetraacetic acid (EDTA). *J. In Vitro Fertil. Embryo Transfer*, **6**, 36–40.
- Kennedy, J. and Donahue, R.P. (1969) Human oocyte: maturation in chemically defined media. *Science*, **164**, 1292–1293.
- Kline, D. and Kline, J.T. (1992) Repetitive calcium transients and the role of calcium in exocytosis and cell cycle activation in the mouse egg. *Dev. Biol.*, **149**, 80–89.
- Kubiak, J., Weber, M. and de Pennart, H. (1993) The metaphase II arrest in mouse oocytes is controlled through microtubule-dependent destruction of cyclin B in the presence of CSF. *EMBO J.*, **12**, 3773–3778.
- Lai, F.A., Erickson, H.P. and Rousseau, E. *et al.* (1988) Purification and reconstitution of the calcium release channel from skeletal muscle. *Nature*, **331**, 315–319.
- Larsen, W.J., Wert, S.E. and Brunner, G.D. (1987) Differential modulation of rat follicle cell gap junction populations at ovulation. *Dev. Biol.*, **122**, 61–71.
- Lauffer, N., Tarlatzis, B.C. and De Cherney, A.H. *et al.* (1984) Asynchrony between human cumulus-corona cell complex and oocyte maturation after human menopausal gonadotropin treatment for *in vitro* fertilization. *Fertil. Steril.*, **42**, 366–369.
- Li, C.C., Chen, E. and O'Connell, C. *et al.* (1993) Detection of c-mos proto-oncogene expression in human cells. *Oncogene*, **8**, 1685–1691.
- Liu, J., Katz, E. and Garcia, J.E. *et al.* (1997) Successful *in vitro* maturation of human oocytes not exposed to human chorionic gonadotropin during ovulation induction, resulting in pregnancy. *Fertil. Steril.*, **67**, 566–568.
- Lorca, T., Cruzalegui, F.H. and Fesquet, D. *et al.* (1993) Calmodulin-dependent protein kinase II mediates inactivation of MPF and CSF upon fertilization of *Xenopus* eggs. *Nature*, **366**, 270–273.
- Ma, S., Kalousek, D.K. and Zouves, C. *et al.* (1989) Chromosome analysis of human oocytes failing to fertilize *in vitro*. *Fertil. Steril.*, **51**, 992–997.
- Macas, E., Floersheim, Y. and Hotz, E. *et al.* (1990) Abnormal chromosomal arrangements in human oocytes. *Hum. Reprod.*, **5**, 703–707.
- Maller, J.L. (1985) Regulation of amphibian oocyte maturation. *Cell Diff.*, **16**, 211–221.
- Mandelbaum, J., Junca, A.M. and Plachot, M. *et al.* (1988) Cryopreservation of human embryos and oocytes. *Hum. Reprod.*, **3**, 117–119.
- Maroulis, G.B., Emery, M. and Verkauf, B.S. *et al.* (1991) Prospective randomized study of human menotropin versus a follicular and a luteal phase gonadotropin-releasing hormone analog–human menotropin stimulation protocols for *in vitro* fertilization. *Fertil. Steril.*, **55**, 1157–1164.
- Martin, R.H., Mahadevan, M.M. and Taylor, P.J. *et al.* (1986) Chromosomal analysis of unfertilized human oocytes. *J. Reprod. Fertil.*, **78**, 673–678.
- Maruo, T., Hayashi, M. and Matsuo, H. *et al.* (1987) The role of thyroid hormone as biological amplifier of the actions of follicle-stimulating hormone in the functional differentiation of cultured porcine granulosa cell. *Endocrinology*, **121**, 1233–1241.
- Maruo, T., Hiramatsu, S. and Otani, T. *et al.* (1992) Increase in the expression of thyroid hormone receptors in porcine granulosa cells early in follicular maturation. *Acta Endocrinol.*, **127**, 152–160.
- Masui, Y. and Clarke, H.J. (1979) Oocyte maturation. *Int. Rev. Cytol.*, **57**, 185–282.
- Mattioli, M. (1992) Biology of oocyte maturation. In Lauria, A. and Gandolfi, F. (eds), *Embryonic Development and Manipulation in Animal Production: Trends in Research and Applications*. London and Chapel, Portland Press, pp. 17–28.
- Mattioli, M., Galeati, G. and Seren, E. (1988) Effect of follicle somatic cells during pig oocyte maturation on egg penetrability and male pronucleus formation. *Gamete Res.*, **20**, 177–183.
- Mehlmann, L.M. and Kline, D. (1994) Regulation of intracellular calcium in the mouse egg: calcium release in response to sperm or inositol trisphosphate is enhanced after meiotic maturation. *Biol. Reprod.*, **51**, 1088–1098.
- Ménézo, Y., Guerin, J. and Czyba, J. (1990) Improvement of human early embryo development *in vitro* by coculture on monolayers of Vero cells. *Biol. Reprod.*, **42**, 301–306.
- Minshall, J., Pines, J. and Golsteyn, R. *et al.* (1989) The role of cyclin synthesis, modification and destruction in the control of cell division. *J. Cell Sci.*, **12** (Suppl.), 77–97.
- Minshall, J., Murray, A. and Colman, A. *et al.* (1991) *Xenopus* oocyte maturation does not require new cyclin synthesis. *J. Cell Biol.*, **114**, 767–772.
- Miyazaki, S., Yazuki, M. and Nakada, K. *et al.* (1992) Block of Ca^{2+} wave and Ca^{2+} oscillations by antibody to the inositol 1,4,5-trisphosphate receptor in fertilized hamster eggs. *Science*, **257**, 251–255.
- Moor, R.M. and Trounson, A.O. (1977) Hormonal and follicular factors affecting maturation of sheep oocytes *in vitro* and their subsequent developmental capacity. *J. Reprod. Fertil.*, **49**, 101–109.
- Morgan, P.M., Warikoo, P.K. and Bavister, B.D. (1991) *In vitro* maturation of oocytes from unstimulated rhesus monkeys: assessment of cytoplasmic maturity by embryonic development after *in vitro* fertilization. *Biol. Reprod.*, **45**, 89–93.
- Mutter, G. and Wolgemuth, D. (1987) Distinct developmental patterns of c-mos proto-oncogene expression in female and male mouse germ cells. *Proc. Natl. Acad. Sci. USA*, **84**, 5301–5305.
- Nagy, Z.P., Cecile, J. and Liu, J. *et al.* (1996) Pregnancy and birth after intracytoplasmic sperm injection of *in vitro* matured germinal-vesicle stage oocytes: case report. *Fertil. Steril.*, **65**, 1047–1050.
- Nakade, S., Rhee, S.K. and Hamaoka, H. *et al.* (1994) Cyclic AMP-dependent phosphorylation of an immunoaffinity purified homotetrameric inositol 1,4,5-trisphosphate receptor (type 1) increases Ca^{2+} flux in reconstituted lipid vesicles. *J. Biol. Chem.*, **269**, 6735–6742.
- Nebreda, A. and Hunter, T. (1993) The c-mos proto-oncogene protein kinase turns on and maintains the activity of MAP kinase, but not MPF in cell free extracts of *Xenopus* oocytes and eggs. *EMBO J.*, **12**, 1979–1983.
- O'Keefe, S.J., Wolfes, H. and Kiessling, A.A. *et al.* (1989) Microinjection of antisense c-mos oligonucleotides prevents meiosis II in the maturing mouse egg. *Proc. Natl. Acad. Sci. USA*, **86**, 7038–7042.
- O'Keefe, S., Kiessling, A. and Cooper, G. (1991) The c-mos gene product is required for cyclin-B accumulation during meiosis of mouse eggs. *Proc. Natl. Acad. Sci. USA*, **88**, 7869–7872.
- Ookata, K., Hisanaga, S. and Okano, T. *et al.* (1992) Relocation and distinct subcellular localization of p34^{cdc2}-cyclin B complex at meiosis reinitiation in starfish oocytes. *EMBO J.*, **11**, 1763–1772.

- Pal, S., Torry, D. and Serta, R. *et al.* (1994) Expression and potential function of the c-mos proto-oncogene in human eggs. *Fertil. Steril.*, **61**, 496–503.
- Palot, M., Harika, G. and Pigeon, F. *et al.* (1988) Propofol in general anesthesia for IVF (by vaginal and transurethral route)—follicular fluid concentration and cleavage rate. *Anesthesiology*, **69**, A573.
- Papadopoulos, G., Randall, J. and Templeton, A.A. (1988) The frequency of chromosome anomalies in human unfertilized oocytes and uncleaved zygotes after insemination *in vitro*. *Hum. Reprod.*, **4**, 568–573.
- Park, S.E., Chung, C.C. and Oum, K.B. *et al.* (1996) Changes in zona pellucida of the human oocytes matured *in vitro* after cryopreservation at the germinal vesicle (GV) stage. *Proceedings of Annual Meeting of American Fertility Society*, Abstr. P-207.
- Park, S.E., Son, W.Y. and Lee, S.H. *et al.* (1997) Chromosome and spindle configurations of human oocytes matured *in vitro* after cryopreservation at the germinal vesicle stage. *Fertil. Steril.*, **68**, 920–926.
- Parkes, A.S. (1958) Factors affecting the viability of frozen ovarian tissue. *J. Endocrinol.*, **17**, 337–343.
- Parrott, D.M.V. (1960) The fertility of mice with orthotopic ovarian grafts derived from frozen tissue. *J. Reprod. Fertil.*, **1**, 230–241.
- Paulson, R.J., Sauer, M.V. and Francis, M.M. *et al.* (1994) Factors affecting pregnancy success of human in-vitro fertilization in unstimulated cycles. *Hum. Reprod.*, **9**, 1571–1575.
- Pellestor, F. (1991) Frequency and distribution of aneuploidy in human female gametes. *Hum. Genet.*, **86**, 283–288.
- Pickering, S.J., Johnson, M.H. and Braude, P.R. *et al.* (1988) Cytoskeletal organization in fresh, aged and spontaneously activated human oocytes. *Hum. Reprod.*, **3**, 978–989.
- Pickering, S.J., Braude, P.R. and Johnson, M.H. *et al.* (1990) Transient cooling to room temperature can cause irreversible disruption of the meiotic spindle in the human oocyte. *Fertil. Steril.*, **54**, 102–108.
- Piko, L. and Clegg, K.B. (1982) Quantitative changes in total RNA, total poly(A), and ribosomes in early mouse embryos. *Dev. Biol.*, **89**, 362–378.
- Pincus, G. and Enzmann, E.V. (1935) The comparative behavior of mammalian eggs *in vivo* and *in vitro*. I. The activation of ovarian eggs. *J. Exp. Med.*, **62**, 655–675.
- Pines, J. and Hunter, T. (1989) Isolation of human cyclin cDNA: evidence for cyclin mRNA and protein regulation in the cell cycle and for interaction with p34^{cdc2}. *Cell*, **58**, 833–846.
- Pines, J. and Hunter, T. (1991) Human cyclins A and B1 are differentially-located in the cell and undergo cell cycle-dependent nuclear transport. *J. Cell Biol.*, **115**, 1–17.
- Piwnicka-Worms, H., Atherton-Fessler, S. and Lee, M. S. *et al.* (1993) Toward an understanding of the eukaryotic cell cycle: a biochemical approach. In Hsueh, A.J.W. and Schomberg, D.W. (eds), *Ovarian Cell Interactions*. Springer-Verlag, New York, pp. 60–78.
- Plachot, M., Veiga, A. and Montagut, J. *et al.* (1988) Are clinical and biological IVF parameters correlated with chromosomal disorders in early life: a multicentric study. *Hum. Reprod.*, **3**, 627–635.
- Prins, G.S., Wagner, C. and Weidel, L. *et al.* (1987) Gonadotropins augment maturation and fertilization of human immature oocytes cultured *in vitro*. *Fertil. Steril.*, **47**, 1035–1037.
- Pyrzynska, B., Maleszewski, M. and Maluchnik, D. (1996) Mouse oocytes penetrated at GV or GVBD stage lose the ability to fuse with additional spermatozoa. *Zygote*, **4**, 123–128.
- Racowsky, C. and Kaufman, M.L. (1992) Nuclear degeneration and meiotic aberration observed in human oocytes matured *in vitro*: analysis by light microscopy. *Fertil. Steril.*, **58**, 750–755.
- Racowsky, C., Prather, A.L. and Johnson, M.K. (1997) Prematurely condensed chromosomes and meiotic abnormalities in unfertilized human oocytes after ovarian stimulation with and without gonadotropin-releasing hormone agonist. *Fertil. Steril.*, **67**, 932–938.
- Regan, L., Owen, E.J. and Jacobs, H.S. (1990) Hypersecretion of luteinising hormone, infertility and miscarriage. *Lancet*, **336**, 1141–1144.
- Richards, J.S. and Hedin, L. (1988) Molecular aspects of hormone action in ovarian follicular development, ovulation, and luteinization. *Annu. Rev. Physiol.*, **50**, 441–463.
- Robertson, J.E. and Baker, R.D. (1969) Role of female sex steroids as possible regulators of oocyte maturation. *Proceedings of the 2nd Annual Meeting of the Society for the Study of Reproduction*, Academic Press, New York. Abstr. 57.
- Rock, J. and Menkin, M.F. (1944) In vitro fertilization and cleavage of human ovarian eggs. *Science*, **100**, 105–107.
- Rodman, T.C. and Bachvarova, R. (1976) RNA synthesis in preovulatory mouse oocytes. *J. Cell Biol.*, **70**, 251–257.
- Roy, L., Singh, B. and Gautier, J. *et al.* (1990) The cyclin component B2 is a substrate for the mos-proto-oncogene product. *Cell*, **61**, 825–831.
- Russell, J.B., Knezevich, K.M. and Fabian, K.F. (1997) Unstimulated immature oocyte retrieval: early versus midfollicular endometrial priming. *Fertil. Steril.*, **67**, 616–620.
- Sagata, N., Oskarsson, M. and Copeland, T. *et al.* (1988) Function of c-mos proto-oncogene product in meiotic maturation in *Xenopus* oocytes. *Nature*, **335**, 519–525.
- Sagata, N., Watanabe, N. and Vande Woude, G.F. *et al.* (1989) The c-mos proto-oncogene product is a cytostatic factor responsible for mitotic arrest in vertebrate eggs. *Nature*, **342**, 512–518.
- Sakkas, D., Batt, P. and Cameron, A. (1989) Development of pre-implantation goat (*Capra hircus*) embryos *in vivo* and *in vitro*. *J. Reprod. Fertil.*, **87**, 359–365.
- Salama, G., Nigam, M. and Shome, K. *et al.* (1992) Ryanodine-affinity chromatography purifies 106 kDa Ca²⁺ release channels from skeletal and cardiac sarcoplasmic reticulum. *Cell Calcium*, **13**, 635–647.
- Santello, L., Alikani, M. and Talansky, B.E. *et al.* (1992) Is the human oocyte plasma membrane polarised? *Hum. Reprod.*, **7**, 999–1003.
- Schroeder, A.C., Downs, S.M. and Eppig, J.J. (1988) Factors affecting the developmental capacity of mouse oocytes undergoing maturation *in vitro*. *Ann. NY Acad. Sci.*, **541**, 197–204.
- Schultz, R.M. and Wassarman, P.M. (1977) Specific changes in the pattern of protein synthesis during meiotic maturation of mammalian oocytes *in vitro*. *Proc. Natl. Acad. Sci. USA*, **74**, 538–541.
- Schultz, R.M., Montgomery, R. and Ward-Bailey, P.F. *et al.* (1983) Regulation of oocyte maturation in the mouse: possible roles of intercellular communication, cAMP and testosterone. *Dev. Biol.*, **95**, 294–304.
- Schultz, G.A., Gifford, D.J. and Mahadevan, M.M. *et al.* (1988) Protein synthetic patterns in immature and mature human oocytes. In Jones, Jr, H.W. and Schrader, C. (eds), *In Vitro Fertilization and Other Assisted Reproduction*. Ann. NY Acad. Sci., **541**, 237–247.
- Shea, B.F., Baker, R.D. and Latour, J.P. (1975) Human follicular oocytes and their maturation *in vitro*. *Fertil. Steril.*, **26**, 1075–1082.
- Shibuya, E.K. and Masui, Y. (1989) Molecular characteristics of cytostatic factors in amphibian egg extracts. *Development*, **106**, 799–808.
- Sirard, M.A. and First, N.L. (1988) In vitro inhibition of oocyte nuclear maturation in the bovine. *Biol. Reprod.*, **39**, 229–234.
- Sirard, M.A., Florman, H.M. and Leibfried-Rutledge, M.L. *et al.* (1989) Timing of nuclear progression and protein synthesis necessary for meiotic maturation in bovine oocytes. *Biol. Reprod.*, **40**, 1257–1263.
- Son, W.Y., Park, S.E. and Lee, K.A. *et al.* (1996) Effects of 1,2-propanediol and freezing-thawing on the *in vitro* developmental capacity of human immature oocytes. *Fertil. Steril.*, **66**, 995–999.
- Sorensen, R.A. and Wassarman, P.M. (1976) Relationship between growth and meiotic maturation of mouse oocytes. *Dev. Biol.*, **50**, 531–536.
- Staigmiller, R.M. and Moor, R.M. (1984) Effect of follicle cells on the maturation and developmental competence of ovine oocytes matured outside the follicle. *Gamete Res.*, **9**, 221–229.
- Stanger, J.D. and Yovich, J.L. (1985) Reduced *in vitro* fertilization of human oocytes from patients with raised basal luteinizing hormone levels during the follicular phase. *Br. J. Obstet. Gynaecol.*, **92**, 385–393.
- Stephoe, P.C. and Edwards, R.G. (1978) Successful birth after IVF. *Lancet*, **ii**, 366.
- Stern, S., Rayyes, A. and Kennedy, J.F. (1972) Incorporation of amino acids during maturation *in vitro* by the mouse oocyte: Effect of puromycin on protein synthesis. *Biol. Reprod.*, **7**, 341–346.
- Sternlicht, A.L. and Schultz, R.M. (1981) Biochemical studies of mammalian oogenesis: kinetics of accumulation of total and poly(A)-containing RNA during growth of the mouse oocyte. *J. Exp. Zool.*, **215**, 191–200.
- Swann, K. (1996) Soluble sperm factors and Ca²⁺ release at fertilization. *Rev. Reprod.*, **1**, 33–39.
- Swann, K. and Lawrence, Y. (1996) How and why spermatozoa cause calcium oscillations in mammalian oocytes. *Mol. Hum. Reprod.*, **2**, 388–390.
- Szybek, K. (1972) In vitro maturation of oocytes from sexually immature mice. *J. Endocrinol.*, **54**, 527–528.

- Tarin, J., Gomez, E. and Sampaio, M. *et al.* (1991) Cytogenetic analysis of human oocytes from fertile women. *Hum. Reprod.*, **6**, 1100–1103.
- Telford, N., Watson, A. and Schultz, G. (1990) Transition from maternal to embryonic control in early mammalian development: A comparison of several species. *Mol. Reprod. Dev.*, **26**, 90–100.
- Tesarik, J. and Sousa, M. (1996) Mechanism of calcium oscillations in human oocytes: a two-store model. *Mol. Hum. Reprod.*, **2**, 383–386.
- Tesarik, J., Kopeny, V. and Plachot, M. *et al.* (1986) Activation of nucleolar and extranuclear RNA synthesis and changes in the ribosomal content of human embryos developing *in vitro*. *J. Reprod. Fertil.*, **78**, 463–470.
- Thibault, C. (1977) Are follicular maturation and oocyte maturation independent processes? *J. Reprod. Fertil.*, **51**, 1–15.
- Thibault, C., Gerard, M. and Ménéz, Y. (1975) Preovulatory and ovulatory mechanisms in oocyte maturation. *J. Reprod. Fertil.*, **45**, 605–610.
- Törmell, J. and Hillensjö, T. (1993) Effect of cyclic AMP on the isolated human oocyte-cumulus complex. *Hum. Reprod.*, **8**, 737–739.
- Toth, T.L., Baka, S.G. and Veeck, L.L. *et al.* (1994a) Fertilization and *in vitro* development of cryopreserved human prophase I oocytes. *Fertil. Steril.*, **61**, 891–894.
- Toth, T.L., Lanzendorf, S.E. and Sandow, B.A. *et al.* (1994b) Cryopreservation of human prophase I oocytes collected from unstimulated follicles. *Fertil. Steril.*, **61**, 1077–1082.
- Trounson, A.O., Mohr, L.R. and Wood, C. *et al.* (1982) Effect of delayed insemination on *in vitro* fertilization, culture and transfer of human embryos. *J. Reprod. Fertil.*, **64**, 285–294.
- Trounson, A., Wood, C. and Kausche, A. (1994) In vitro maturation and fertilization and developmental competence of oocytes recovered from untreated polycystic ovarian patients. *Fertil. Steril.*, **62**, 353–362.
- Tsuji, K., Sowa, M. and Nakano, R. (1985) Relationship between human oocyte maturation and different follicular sizes. *Biol. Reprod.*, **32**, 413–417.
- Van Blerkom, J. (1991) Microtubule mediation of cytoplasmic and nuclear maturation during the early stages of resumed meiosis in cultured mouse oocytes. *Proc. Natl. Acad. Sci. USA*, **88**, 5031–5035.
- Van Blerkom, J. (1994) Developmental failure in human reproduction associated with chromosomal abnormalities and cytoplasmic pathologies in meiotically-mature human oocytes. In Van Blerkom, J. (ed.), *The Biological Basis of Early Reproductive Failure in the Human: Applications to Medically Assisted Conception*. Oxford University Press, Oxford, pp. 283–325.
- Van Blerkom, J. (1996) The influence of intrinsic and extrinsic factors on the developmental potential of chromosomal normality of the human oocyte. *J. Soc. Gyn. Invest.*, **3**, 3–11.
- Van Blerkom, J. and Henry, G. (1992) Oocyte dysmorphism and aneuploidy in meiotically-mature human oocytes after ovarian stimulation. *Hum. Reprod.*, **7**, 379–390.
- Van Blerkom, J. and Davis, P. (1994) Cytogenetic, cellular and developmental consequences of cryopreservation of immature and mature mouse and human oocytes. *Microsc. Res. Techn.*, **27**, 165–193.
- Van Blerkom, J., Davis, P. and Lee, J. (1995) ATP content of human oocytes and developmental potential and outcome after *in vitro* fertilization and embryo transfer. *Hum. Reprod.*, **10**, 415–424.
- Vanderhyden, B.C. and Armstrong, D.T. (1989) Role of cumulus cells and serum on the *in vitro* maturation, fertilization, and subsequent development of rat oocytes. *Biol. Reprod.*, **40**, 720–728.
- Van der Elst, J., Van den Abbeel, E. and Jacobs, R. *et al.* (1988) Effect of 1,2-propanediol and dimethyl sulfoxide on the meiotic spindle of the mouse oocyte. *Hum. Reprod.*, **8**, 960–967.
- Van Uem, J.F.H.M., Siebzehnriuel, E.R. and Schuh, B. *et al.* (1987) Birth after cryopreservation of unfertilized oocytes. *Lancet*, **i**, 752–753.
- Veeck, L.L., Wortham, J.W.E. and Witmyer, J. *et al.* (1983) Maturation and fertilization of morphologically immature human oocytes in a program of *in vitro* fertilization. *Fertil. Steril.*, **39**, 594–602.
- Venturoli, S., Orsini, L.F. and Paradisi, R. *et al.* (1986) Human urinary follicle stimulating hormone and menopausal gonadotropin in induction of multiple follicle growth and ovulation. *Fertil. Steril.*, **45**, 30–35.
- Vincent, C. and Johnson, M.H. (1992) Cooling, cryoprotectants, and the cytoskeleton of the mammalian oocytes. *Oxf. Rev. Reprod. Biol.*, **14**, 73–100.
- Wall, M., Marks, K. and Smith, T. *et al.* (1996) Cytogenetic and fluorescence *in situ* hybridization chromosome studies on *in vitro* fertilized and intracytoplasmic sperm injected 'failed-fertilized' human oocytes. *Hum. Reprod.*, **11**, 2230–2238.
- Wassarman, P. and Kinloch, R. (1992) Gene expression during oogenesis in mice. *Mutat. Res.*, **296**, 3–15.
- Wassarman, P.M. and Letourneau, G.E. (1976) RNA synthesis in fully grown mouse oocytes. *Nature*, **361**, 73–74.
- Watanabe, N., Vande Woude, G.F. and Ikawa, Y. *et al.* (1989) Specific proteolysis of the c-mos oncogene by calpain on fertilization of *Xenopus* eggs. *Nature*, **342**, 505–511.
- Weston, A.M., Zelinski-Wooten, M.B. and Hutchison, J.S. *et al.* (1996) Developmental potential of embryos produced by in-vitro fertilization from gonadotrophin-releasing hormone antagonist-treated macaques stimulated with recombinant human follicle stimulating hormone alone or in combination with luteinizing hormone. *Hum. Reprod.*, **11**, 608–613.
- Whitaker, M. (1996) Control of meiotic arrest. *Rev. Reprod.*, **1**, 127–135.
- Whitaker, M. and Swann, K. (1993) Lighting the fuse at fertilization. *Development*, **117**, 1–12.
- Wickramasinghe, D., Ebert, K.M. and Albertini, D.F. (1991) Meiotic competence acquisition is associated with the appearance of M-phase characteristics in growing mouse oocytes. *Dev. Biol.*, **143**, 162–172.
- Wramsby, H., Liedholm, P. (1984) A gradual fixation method for chromosomal preparation of human oocytes. *Fertil. Steril.*, **41**, 736–738.
- Wu, S. and Wolgemuth, D.J. (1995) The distinct and developmentally regulated patterns of expression of members of the mouse Cdc25 gene family suggest differential functions during gametogenesis. *Dev. Biol.*, **170**, 195–206.
- Wu, T.C., Wang, L. and Wan, Y.J. (1993) Detection of estrogen receptor messenger ribonucleic acid in human oocytes and cumulus-oocyte complexes using reverse transcriptase-polymerase chain reaction. *Fertil. Steril.*, **59**, 54–59.
- Zhang, X., Zerafa, A. and Wong, J. *et al.* (1993) Human menopausal gonadotropin during *in vitro* maturation of human oocytes retrieved from small follicles enhances *in vitro* fertilization and cleavage rates. *Fertil. Steril.*, **59**, 850–853.
- Zhang, S.S., Carrillo, A.J. and Darling, D.S. (1997) Expression of multiple thyroid hormone receptor mRNAs in human oocytes, cumulus cells, and granulosa cells. *Mol. Hum. Reprod.*, **3**, 555–562.

Received on July 31, 1997; accepted on March 9, 1998